

A REACTION COEFFICIENT IDENTIFICATION PROBLEM FOR FRACTIONAL DIFFUSION*

ENRIQUE OTÁROLA[†] AND TRAN NHAN TAM QUYEN[‡]

Abstract. We analyze a coefficient identification problem for the spectral fractional powers of a symmetric, coercive, linear, elliptic, second-order operator in a bounded domain Ω . We realize fractional diffusion as the Dirichlet-to-Neumann map for a nonuniformly elliptic problem posed on the semi-infinite cylinder $\Omega \times (0, \infty)$. We thus consider an equivalent coefficient identification problem, where the coefficient to be identified appears explicitly. We derive existence of local solutions, optimality conditions and a rapid decay of solutions on the extended domain $(0, \infty)$. The latter property suggests a truncation that is suitable for numerical approximation. We thus propose and analyze a fully discrete scheme that discretizes the set of admissible coefficients with piecewise constant functions. The discretization of the state equation relies on the tensorization of a first-degree FEM in Ω with a suitable hp -FEM in the extended dimension. We derive convergence results and obtain a priori error estimates.

Key words. coefficient identification problems, fractional diffusion, nonlocal operators, finite elements, error estimates.

AMS subject classifications. 35J70, 35R11, 35R30.

1. Introduction. In this work we shall be interested in the analysis of a coefficient identification problem for certain fractional powers of symmetric, coercive, self-adjoint, second order differential operators in bounded domains with homogeneous Dirichlet boundary conditions. To make matters precise, for $d \geq 1$, we let $\Omega \subset \mathbb{R}^d$ be an open and bounded domain with Lipschitz boundary $\partial\Omega$. We define

$$\mathcal{L}w := -\operatorname{div}(A\nabla w) + qw, \tag{1.1}$$

supplemented with homogeneous Dirichlet boundary conditions. The diffusion matrix $A \in L^\infty(\Omega, \operatorname{GL}(\mathbb{R}^d))$ is symmetric and uniformly positive definite and the reaction term q belongs to the following set of admissible coefficients:

$$\mathcal{Q} := \{q \in L^\infty(\Omega) \mid 0 \leq q(x) \leq \bar{q} \text{ a.e. in } \Omega\}, \quad \bar{q} > 0. \tag{1.2}$$

For $s \in (0, 1)$, we denote by \mathcal{L}^s the spectral fractional powers of the operator \mathcal{L} .

Given $s \in (0, 1)$ and f , we shall be concerned with the analysis of the problem of identifying the coefficient q in the Dirichlet problem for fractional diffusion

$$\mathcal{L}^s u = f \text{ in } \Omega \tag{1.3}$$

from the observation data $z_\delta \in L^2(\Omega)$ of the exact solution u^\dagger in the domain Ω . The aforementioned data satisfies that

$$\|u^\dagger - z_\delta\|_{L^2(\Omega)} \leq \delta, \tag{1.4}$$

* The research of EO was supported in part by CONICYT through project FONDECYT 3160201. TNTQ gratefully acknowledges support of the University of Goettingen, Goettingen, Lower Saxony, Germany.

[†]Departamento de Matemática, Universidad Técnica Federico Santa María, Valparaíso, Chile (enrique.otarola@usm.cl).

[‡]Institute for Numerical and Applied Mathematics, University of Goettingen, Goettingen, Germany (quyen.tran@uni-goettingen.de).

where $\delta > 0$ denotes the level of measurement error. To solve the proposed identification problem in a stable manner, we will utilize the standard output least squares method with Tikhonov regularization [16, 20, 40]. In fact, for estimating the coefficient q in (1.3) from the observations z_δ of the exact solution u^\dagger , we will invoke the following cost functional

$$J_{\delta,\rho}(q) := \frac{1}{2} \|U(q) - z_\delta\|_{L^2(\Omega)}^2 + \frac{\rho}{2} \|q - q^*\|_{L^2(\Omega)}^2, \quad (1.5)$$

where U denotes the so-called coefficient-to-solution operator, which associates to an element $q \in \mathcal{Q}$ a unique weak solution $u =: U(q)$ of problem (1.3). In (1.5), q^* corresponds to an a priori estimate of the coefficient q to be identified and $\rho > 0$ denotes the so-called regularization parameter. We will thus consider a minimizer of the optimization problem

$$\min_{q \in \mathcal{Q}} J_{\delta,\rho}(q) \quad (1.6)$$

as a reconstruction. One of the main difficulties in the analysis and design of solution techniques for problem (1.6) is that \mathcal{L}^s is a *nonlocal* operator [8, 9, 10, 11, 13, 14]. We must immediately notice that the coefficient q to be identified does not appear explicitly neither in the strong nor the weak formulation of problem (1.3). The coefficient q acts, however, modifying the spectrum of the differential operator \mathcal{L} and thus the definition of the fractional powers \mathcal{L}^s ; see definition (2.4) below. As a consequence, the analysis of the coefficient identification problem (1.6) is intricate.

The mathematical analysis of fractional diffusion has been one of the most studied topics in the past decade [8, 9, 10, 11, 12, 14, 34, 46]. A breakthrough in the theory, that allows for the localization of \mathcal{L}^s , is due to Caffarelli and Silvestre [11]. When $\Omega = \mathbb{R}^d$ and $\mathcal{L} = -\Delta$, i.e., in the case when \mathcal{L} coincides with the Laplace operator in \mathbb{R}^d , Caffarelli and Silvestre proved that any power $s \in (0, 1)$ of the fractional Laplacian $(-\Delta)^s$ can be realized as the Dirichlet-to-Neumann map for an extension problem posed on the upper half-space $\mathbb{R}_+^{d+1} := \mathbb{R}^d \times (0, \infty)$ [11]. Such an extension problem involves a nonuniformly but *local* elliptic PDE. This result was later adapted in [10, 14, 42] to bounded domains Ω and more general operators, thereby obtaining an extension problem posed on the semi-infinite cylinder $\mathcal{C} := \Omega \times (0, \infty)$. The latter corresponds to the following *local* boundary value problem

$$\begin{cases} -\operatorname{div}(y^\alpha \mathbf{A} \nabla \mathcal{U}) + q y^\alpha \mathcal{U} = 0 & \text{in } \mathcal{C}, \\ \mathcal{U} = 0 & \text{on } \partial_L \mathcal{C}, \\ \partial_{\nu^\alpha} \mathcal{U} = d_s f & \text{on } \Omega \times \{0\}, \end{cases} \quad (1.7)$$

where $\mathbf{A} = \operatorname{diag}\{A, 1\} \in L^\infty(\mathcal{C}, \operatorname{GL}(\mathbb{R}^{d+1}))$ and $\partial_L \mathcal{C} := \partial\Omega \times (0, \infty)$ denotes the lateral boundary of \mathcal{C} . In addition, in (1.7), d_s denotes a positive normalization constant given by $d_s := 2^{1-2s} \Gamma(1-s)/\Gamma(s)$ and the parameter α is defined as $\alpha := 1 - 2s \in (-1, 1)$ (cf. [11, 42]). The conormal exterior derivative of \mathcal{U} at $\Omega \times \{0\}$ is defined by

$$\partial_{\nu^\alpha} \mathcal{U} = - \lim_{y \rightarrow 0^+} y^\alpha \mathcal{U}_y, \quad (1.8)$$

where the limit is understood in the distributional sense. We will call $y \in (0, \infty)$ the *extended variable* and call the dimension $d+1$ in \mathbb{R}_+^{d+1} the *extended dimension*. With the extension \mathcal{U} at hand, the fractional powers of \mathcal{L} in (1.3) and the Dirichlet-to-Neumann operator of problem (1.7) are related by

$$d_s \mathcal{L}^s u = \partial_{\nu^\alpha} \mathcal{U} \text{ in } \Omega. \quad (1.9)$$

Motivated by applications to tomography and related techniques, the study of parameter identification problems in partial differential equations (PDEs) has received considerable attention over the past 50 years. A rather incomplete list of problems where parameter identification problems appear includes modern medical imaging modalities, aquifer analysis, geophysical prospecting and pollutant detection. We refer the interested reader to [3, 4, 18, 41, 43] for a survey. In particular, the problem of identifying the reaction coefficient in local and elliptic equations has been extensively studied; we refer the reader to [27, 28] and references therein. In contrast to these advances, and to the best of our knowledge, this is the first work addressing the study of a reaction coefficient identification problem for fractional diffusion.

We provide a comprehensive treatment for a reaction coefficient identification problem for the spectral fractional powers of a symmetric, coercive, linear, elliptic, second-order operator in a bounded domain Ω . We overcome the nonlocality of fractional diffusion by using the results of Caffarelli and Silvestre [11]. We realize (1.3) by (1.7) and propose an equivalent identification problem. As a consequence, standard variational techniques are applicable since, in contrast to (1.3), the reaction coefficient q appears explicitly in the weak formulation of (1.7). This is one of the highlights of our work. We rigorously derive existence and differentiability results, optimality conditions and regularity estimates. We also present a numerical scheme, and prove that there exists a sequence of local minima that converges to a local and exact solution for all values of s .

Our presentation is organized as follows. The notation and functional setting is described in section 2, where we also briefly describe, in §2.1, the definition of spectral fractional diffusion and, in §2.2, its localization via the Caffarelli–Silvestre extension property. In section 3 we introduce the *extended* identification problem and prove that is equivalent to (1.6). In addition, we derive the existence of local solutions, study differentiability properties for the underlying coefficient-to-solution operator, analyze optimality conditions and derive regularity estimates. In section 4, we begin the numerical analysis of our problem. We introduce a truncation of the state equation and a *truncated* identification problem. We derive approximation properties of its solution. In section 5.1 we briefly recall the finite element scheme of [2] that approximates the solution to (1.7). In section 5.2 we introduce a fully discrete scheme for the truncated identification problem and derive convergence of discrete solutions when the regularization parameter converges to zero. Finally, in section 5.3 we derive convergence results and a priori error estimates for the proposed fully discrete scheme.

2. Notation and preliminaries. We adopt the notation of [36, 37]. Throughout this work Ω is a convex polytopal subset of \mathbb{R}^d , $d \geq 1$, with boundary $\partial\Omega$. Besides the semi-infinite cylinder $\mathcal{C} = \Omega \times (0, \infty)$, we introduce the truncated cylinder with base Ω and height \mathcal{Y} as $\mathcal{C}_{\mathcal{Y}} := \Omega \times (0, \mathcal{Y})$, and its lateral boundary $\partial_L \mathcal{C} := \partial\Omega \times (0, \mathcal{Y})$. Since the extended variable y will play a special role in the analysis that we will perform, throughout the text, points $x \in \mathcal{C} = \Omega \times (0, \infty) \subset \mathbb{R}^{d+1}$ will be written as

$$x = (x', y), \quad x' \in \Omega \subset \mathbb{R}^d, \quad y \in (0, \infty).$$

Whenever X is a normed space we denote by $\|\cdot\|_X$ its norm and by X' its dual. For normed spaces X and Y we write $X \hookrightarrow Y$ to indicate continuous embedding.

By $a \lesssim b$ we mean $a \leq Cb$ with a constant C that neither depends on a , b nor the discretization parameters. The notation $a \sim b$ signifies $a \lesssim b \lesssim a$. The value of C might change at each occurrence.

Finally, since we assume Ω to be convex, in what follows we will make use, without explicit mention, of the following regularity result [26]:

$$\|w\|_{H^2(\Omega)} \lesssim \|\mathcal{L}w\|_{L^2(\Omega)} \quad \forall w \in H^2(\Omega) \cap H_0^1(\Omega). \quad (2.1)$$

2.1. Fractional powers of elliptic operators. In this section, we invoke spectral theory [5, 29] and define the *spectral fractional powers* of the elliptic operator \mathcal{L} . To accomplish this task, we begin by noticing that in view of the assumptions on A and q , the operator \mathcal{L} induces the following inner product on $H_0^1(\Omega)$:

$$a_\Omega : H_0^1(\Omega) \times H_0^1(\Omega) \rightarrow \mathbb{R}, \quad a_\Omega(w, v) = \int_\Omega (A\nabla w \cdot \nabla v + q w v) \, dx'. \quad (2.2)$$

In addition, $\mathcal{L} : H_0^1(\Omega) \rightarrow H^{-1}(\Omega)$ defined by $u \mapsto a_\Omega(u, \cdot)$ is an isomorphism. The eigenvalue problem

$$(\lambda, \phi) \in \mathbb{R} \times H_0^1(\Omega) \setminus \{0\} : \quad a_\Omega(\phi, v) = \lambda(\phi, v)_{L^2(\Omega)} \quad \forall v \in H_0^1(\Omega)$$

has a countable collection of solutions $\{\lambda_k, \varphi_k\}_{k \in \mathbb{N}} \subset \mathbb{R}^+ \times H_0^1(\Omega)$, where $\{\varphi_k\}_{k \in \mathbb{N}}$ is an orthonormal basis of $L^2(\Omega)$ and an orthogonal basis of $(H_0^1(\Omega), a_\Omega(\cdot, \cdot))$. The real sequence of eigenvalues $\{\lambda_k\}_{k \in \mathbb{N}}$ is enumerated in increasing order, counting multiplicities and it accumulates at infinite.

With these ingredients at hand, we define, for $s \geq 0$, the fractional Sobolev space

$$\mathbb{H}^s(\Omega) = \left\{ w = \sum_{k=1}^{\infty} w_k \varphi_k \mid \|w\|_{\mathbb{H}^s(\Omega)} := \left(\sum_{k=1}^{\infty} \lambda_k^s w_k^2 < \infty \right)^{\frac{1}{2}} \right\}. \quad (2.3)$$

We must immediately mention that in view of the results of [29] the space $\mathbb{H}^s(\Omega)$ can be equivalently defined in terms of the eigenpairs $\{\mu_k, \phi_k\}_{k \in \mathbb{N}}$ of the Dirichlet Laplace operator in the bounded domain Ω .

We denote by $\langle \cdot, \cdot \rangle$ the duality pairing between $\mathbb{H}^s(\Omega)$ and $\mathbb{H}^{-s}(\Omega)$; $\mathbb{H}^{-s}(\Omega)$ denoting the dual space of $\mathbb{H}^s(\Omega)$. Through this duality pairing, the definition of the space (2.3) and the norm $\|\cdot\|_{\mathbb{H}^s(\Omega)}$ can both be extended to $s < 0$. In addition, by real interpolation between $L^2(\Omega)$ and $H_0^1(\Omega)$, we infer, for $s \in (0, 1)$, that $\mathbb{H}^s(\Omega) = [L^2(\Omega), H_0^1(\Omega)]_s$. If $s \in (1, 2]$, owing to (2.1), we have that $\mathbb{H}^s(\Omega) = H^s(\Omega) \cap H_0^1(\Omega)$ [24].

For $s = 1$ and $w = \sum_{k=1}^{\infty} w_k \varphi_k \in \mathbb{H}^1(\Omega)$, we thus have that $\mathcal{L}w = \sum_{k=1}^{\infty} \lambda_k w_k \varphi_k \in \mathbb{H}^{-1}(\Omega)$. For $s \in (0, 1)$ and $w = \sum_{k=1}^{\infty} w_k \varphi_k \in \mathbb{H}^s(\Omega)$, the operator \mathcal{L}^s is defined by

$$\mathcal{L}^s : \mathbb{H}^s(\Omega) \rightarrow \mathbb{H}^{-s}(\Omega), \quad \mathcal{L}^s w = \sum_{k=1}^{\infty} \lambda_k^s w_k \varphi_k. \quad (2.4)$$

A weak formulation for (1.3) reads as follows: Find $u \in \mathbb{H}^s(\Omega)$ such that

$$\langle \mathcal{L}^s u, v \rangle = \langle f, v \rangle \quad \forall v \in \mathbb{H}^s(\Omega). \quad (2.5)$$

Given $f \in \mathbb{H}^{-s}(\Omega)$, problem (2.5) admits a unique weak solution $u \in \mathbb{H}^s(\Omega)$ [10, 14]. In addition, the following estimate can be derived [10, 14]

$$\|u\|_{\mathbb{H}^s(\Omega)} \lesssim \|f\|_{\mathbb{H}^{-s}(\Omega)}. \quad (2.6)$$

2.2. An extension property. The Caffarelli–Silvestre extension result requires us to deal with the local but nonuniformly elliptic problem (1.7) (cf. [6, 10, 11, 14]). It is thus instrumental to define Lebesgue and Sobolev spaces with the weight y^α , where $\alpha \in (-1, 1)$. If $D \subset \mathbb{R}^{d+1}$, we define $L^2(y^\alpha, D)$ as the Lebesgue space for the measure $|y|^\alpha dx$. We also define the weighted Sobolev space

$$H^1(y^\alpha, D) := \{w \in L^2(y^\alpha, D) \mid |\nabla w| \in L^2(y^\alpha, D)\},$$

where ∇w is the distributional gradient of w . We equip $H^1(y^\alpha, D)$ with the norm

$$\|w\|_{H^1(y^\alpha, D)} = \left(\|w\|_{L^2(y^\alpha, D)}^2 + \|\nabla w\|_{L^2(y^\alpha, D)}^2 \right)^{\frac{1}{2}}. \quad (2.7)$$

Since $\alpha \in (-1, 1)$, the weight y^α belongs to the Muckenhoupt class $A_2(\mathbb{R}^{d+1})$ (cf. [19, 22, 25, 35, 45]). This in particular implies that $H^1(y^\alpha, D)$ with norm (2.7) is a Hilbert space and $C^\infty(D) \cap H^1(y^\alpha, D)$ is dense in $H^1(y^\alpha, D)$ (cf. [45, Proposition 2.1.2, Corollary 2.1.6], [32] and [25, Theorem 1]).

We define the weighted Sobolev space

$$\mathring{H}^1(y^\alpha, \mathcal{C}) = \{w \in H^1(y^\alpha, \mathcal{C}) \mid w = 0 \text{ on } \partial_L \mathcal{C}\}, \quad (2.8)$$

and immediately notice that $\mathring{H}^1(y^\alpha, \mathcal{C})$ can be equivalently defined as [14]

$$\begin{aligned} \mathring{H}^1(y^\alpha, \mathcal{C}) = \{w : \mathcal{C} \rightarrow \mathbb{R} : & \quad w \in H^1(\Omega \times (r, t)) \quad \forall 0 < r < t < \infty, \\ & \quad w = 0 \text{ on } \partial_L \mathcal{C}, \quad \|\nabla w\|_{L^2(y^\alpha, \mathcal{C})} < \infty\}. \end{aligned} \quad (2.9)$$

As [36, inequality (2.21)] shows, the following *weighted Poincaré inequality* holds:

$$\|w\|_{L^2(y^\alpha, \mathcal{C})} \lesssim \|\nabla w\|_{L^2(y^\alpha, \mathcal{C})} \quad \forall w \in \mathring{H}^1(y^\alpha, \mathcal{C}). \quad (2.10)$$

Consequently, the seminorm on $\mathring{H}^1(y^\alpha, \mathcal{C})$ is equivalent to (2.7).

For $w \in H^1(y^\alpha, \mathcal{C})$, $\text{tr}_\Omega w$ denotes its trace onto $\Omega \times \{0\}$ which satisfies

$$\text{tr}_\Omega \mathring{H}^1(y^\alpha, \mathcal{C}) = \mathbb{H}^s(\Omega), \quad \|\text{tr}_\Omega w\|_{\mathbb{H}^s(\Omega)} \leq C_{\text{tr}_\Omega} \|w\|_{\mathring{H}^1(y^\alpha, \mathcal{C})}; \quad (2.11)$$

see [36, Proposition 2.5]. We notice that, if a function w belongs to $\mathring{H}^1(y^\alpha, \mathcal{C})$ then, in view of (2.9), we have, for $y > 0$, that $w(\cdot, y) \in H^{1/2}(\Omega)$.

Define the continuous and coercive bilinear form $a_{\mathcal{C}} : \mathring{H}^1(y^\alpha, \mathcal{C}) \times \mathring{H}^1(y^\alpha, \mathcal{C}) \rightarrow \mathbb{R}$:

$$a_{\mathcal{C}}(w, \phi)(q) = \int_{\mathcal{C}} y^\alpha (\mathbf{A} \nabla w \cdot \nabla \phi + qw\phi) \, dx' \, dy. \quad (2.12)$$

We shall simply write $a_{\mathcal{C}}(w, \phi)$ or $a_{\mathcal{C}}$ when no confusion arises. Notice that $a_{\mathcal{C}}$ induces an inner product on $\mathring{H}^1(y^\alpha, \mathcal{C})$ and an *energy norm*. The latter is defined as follows:

$$\|w\|_{\mathcal{C}} := a_{\mathcal{C}}(w, w)^{1/2} \quad \forall w \in \mathring{H}^1(y^\alpha, \mathcal{C}). \quad (2.13)$$

In view of the assumptions on \mathbf{A} and c , we conclude that $\|w\|_{\mathcal{C}} \sim \|\nabla w\|_{L^2(y^\alpha, \mathcal{C})}$.

We now present a weak formulation for problem (1.7):

$$\mathcal{W} \in \mathring{H}^1(y^\alpha, \mathcal{C}) : \quad a_{\mathcal{C}}(\mathcal{W}, \phi)(q) = d_s \langle f, \text{tr}_\Omega \phi \rangle \quad \forall \phi \in \mathring{H}^1(y^\alpha, \mathcal{C}). \quad (2.14)$$

The fundamental result of Caffarelli and Silvestre [11], [10, Proposition 2.2], [14], [42, Theorem 1.1] is stated below.

PROPOSITION 2.1 (Caffarelli–Silvestre extension result). *Let $s \in (0, 1)$ and $u \in \mathbb{H}^s(\Omega)$ be the solution to (1.3) with $f \in \mathbb{H}^{-s}(\Omega)$. If \mathcal{U} solves (2.14), then*

$$u = \text{tr}_\Omega \mathcal{U} \text{ in } \Omega, \quad d_s \mathcal{L}^s u = \partial_{\nu^\alpha} \mathcal{U} \text{ in } \Omega, \quad (2.15)$$

where $d_s = 2^{1-2s} \Gamma(1-s) / \Gamma(s)$.

The $\mathbb{H}^s(\Omega)$ -norm of u and the energy norm $\|\cdot\|_{\mathcal{C}}$ of \mathcal{U} are related by

$$d_s \|u\|_{\mathbb{H}^s(\Omega)}^2 = \|\mathcal{U}\|_{\mathcal{C}}^2. \quad (2.16)$$

We conclude this section by proving a representation formula for the solution \mathcal{U} of problem (2.14). If $u = \sum_{k=1}^{\infty} u_k \varphi_k$ solves (1.3), then the unique solution \mathcal{U} of problem (2.14) admits the representation [36, formula (2.24)]

$$\mathcal{U}(x', y) = \sum_{k=1}^{\infty} u_k \varphi_k(x') \psi_k(y), \quad u_k := \lambda_k^{-s} f_k, \quad f_k = (f, \varphi_k)_{L^2(\Omega)}. \quad (2.17)$$

We recall that $\{\lambda_k, \varphi_k\}_{k \in \mathbb{N}}$ is the set of eigenpairs of \mathcal{L} . The functions ψ_k solve

$$\begin{cases} \psi_k''(y) + \alpha y^{-1} \psi_k'(y) - \lambda_k \psi_k(y) = 0, & y \in (0, \infty), \\ \psi_k(0) = 1, & \lim_{y \rightarrow \infty} \psi_k(y) = 0. \end{cases} \quad (2.18)$$

Thus, if $s = \frac{1}{2}$, we have $\psi_k(y) = \exp(-\sqrt{\lambda_k} y)$ [10, Lemma 2.10]; more generally, if $s \in (0, 1) \setminus \{\frac{1}{2}\}$, then [14, Proposition 2.1]

$$\psi_k(y) = c_s (\sqrt{\lambda_k} y)^s K_s(\sqrt{\lambda_k} y),$$

where $c_s = 2^{1-s} / \Gamma(s)$ and K_s denotes the modified Bessel function of the second kind. We refer the reader to [1, Chapter 9.6] for a comprehensive treatment of the Bessel function K_s .

3. The extended identification problem. In order to analyze problem (1.6) and design a numerical technique to efficiently solve it, we will consider an equivalent minimization problem based on (2.14): the *extended identification problem*. To describe it, we define the extended coefficient-to-solution operator

$$E : \mathcal{Q} \rightarrow \mathring{H}^1(y^\alpha, \mathcal{C}), \quad q \mapsto \mathcal{U}(q), \quad (3.1)$$

which, for given a coefficient $q \in \mathcal{Q}$ associates to it the unique weak solution $\mathcal{U} =: E(q) \in \mathring{H}^1(y^\alpha, \mathcal{C})$ of problem (2.14). With the map E at hand, we define, for $\rho > 0$ and $z_\delta \in L^2(\Omega)$, the cost functional

$$\mathcal{J}_{\delta, \rho}(q) := \frac{1}{2} \|\text{tr}_\Omega E(q) - z_\delta\|_{L^2(\Omega)}^2 + \frac{\rho}{2} \|q - q^*\|_{L^2(\Omega)}^2. \quad (3.2)$$

The *extended identification problem* thus reads as follows:

$$\min_{q \in \mathcal{Q}} \mathcal{J}_{\delta, \rho}(q) \quad (3.3)$$

In the following result we state the equivalence between the fractional identification problem (1.6) and the extended one (3.3).

THEOREM 3.1 (equivalence of (1.6) and (3.3)). If $(E(q^*), q^*) \in \mathring{H}^1(y^\alpha, \mathcal{C}) \times \mathcal{Q}$ and $(U(q^\dagger), q^\dagger) \in \mathbb{H}^s(\Omega) \times \mathcal{Q}$ solve (3.3) and (1.6), respectively, then

$$q^* = q^\dagger, \quad \text{tr}_\Omega E(q^*) = U(q^*), \quad (3.4)$$

where tr_Ω is defined as in §2.2.

Proof. The proof is a direct consequence of the identity (2.15). \square

3.1. Existence of solutions. We present the following result.

THEOREM 3.2 (existence of solutions). *The regularization problem (3.3) has a solution $q_{\delta, \rho}$.*

Proof. Let $(q_n)_{n \in \mathbb{N}} \subset \mathcal{Q}$ be a minimizing sequence for problem (3.3), i.e.,

$$\begin{aligned} & \lim_{n \rightarrow \infty} \left(\frac{1}{2} \|\text{tr}_\Omega E(q_n) - z_\delta\|_{L^2(\Omega)}^2 + \frac{\rho}{2} \|q_n - q^*\|_{L^2(\Omega)}^2 \right) \\ &= \inf_{q \in \mathcal{Q}} \left(\frac{1}{2} \|\text{tr}_\Omega E(q) - z_\delta\|_{L^2(\Omega)}^2 + \frac{\rho}{2} \|q - q^*\|_{L^2(\Omega)}^2 \right); \end{aligned}$$

such a sequence does exist by the definition of the infimum. We immediately notice that in view of [39, Theorem 1.7] we can conclude that \mathcal{Q} is a weakly* compact subset of $L^\infty(\Omega)$; see also [38, Remark 2.1] and [7, Theorem 3.16]. Consequently, we deduce the existence of $q_{\delta, \rho} \in \mathcal{Q}$ and a subsequence $(q_{n_k})_{k \in \mathbb{N}} \subset (q_n)_{n \in \mathbb{N}}$ such that $(q_{n_k})_{k \in \mathbb{N}}$ converges weakly* to $q_{\delta, \rho}$ in $L^\infty(\Omega)$, i.e.,

$$\lim_{k \rightarrow \infty} \int_\Omega q_{n_k} \chi \, dx' = \int_\Omega q_{\delta, \rho} \chi \, dx' \quad \forall \chi \in L^1(\Omega). \quad (3.5)$$

On the other hand, in view of (2.6) and (2.16), we have that $(E_{n_k})_{n \in \mathbb{N}}$, defined by $E_{n_k} := E(q_{n_k})$, is bounded in $H^1(y^\alpha, \mathcal{C})$. Consequently, we conclude the existence of $\Theta \in H^1(y^\alpha, \mathcal{C})$ and a nonrelabeled subsequence $(E_{n_k})_{k \in \mathbb{N}}$ such that

$$E_{n_k} \rightharpoonup \Theta \text{ in } H^1(y^\alpha, \mathcal{C}), \quad E_{n_k} \rightharpoonup \Theta \text{ in } L^2(y^\alpha, \mathcal{C}), \quad k \uparrow \infty. \quad (3.6)$$

In addition, since $\text{tr}_\Omega : \mathring{H}^1(y^\alpha, \mathcal{C}) \rightarrow \mathbb{H}^s(\Omega)$ is linear and continuous, the compact embedding $\mathbb{H}^s(\Omega) \hookrightarrow L^2(\Omega)$ [33, Theorem 3.27] reveals that

$$\text{tr}_\Omega E_{n_k} \rightharpoonup \text{tr}_\Omega \Theta \text{ in } \mathbb{H}^s(\Omega), \quad \text{tr}_\Omega E_{n_k} \rightarrow \text{tr}_\Omega \Theta \text{ in } L^2(\Omega), \quad k \uparrow \infty. \quad (3.7)$$

In what follows we derive that $\Theta = E(q_{\delta, \rho})$. Let $n_k \in \mathbb{N}$ and $\phi \in \mathring{H}^1(y^\alpha, \mathcal{C})$. From definitions (2.14) and (2.12), we obtain that

$$\begin{aligned} d_s \langle f, \text{tr}_\Omega \phi \rangle &= \int_{\mathcal{C}} y^\alpha (\mathbf{A} \nabla E_{n_k} \cdot \nabla \phi + q_{n_k} E_{n_k} \phi) \, dx = \int_{\mathcal{C}} y^\alpha (q_{n_k} - q_{\delta, \rho}) \Theta \phi \, dx \\ &+ \int_{\mathcal{C}} y^\alpha (\mathbf{A} \nabla E_{n_k} \cdot \nabla \phi + q_{\delta, \rho} \Theta \phi) \, dx + \int_{\mathcal{C}} y^\alpha q_{n_k} (E_{n_k} - \Theta) \phi \, dx =: \text{I}_k + \text{II}_k + \text{III}_k. \end{aligned}$$

We proceed to estimate I_k . To accomplish this task, we define, a.e. $x' \in \Omega$, the function $\chi(x') := \int_0^\infty y^\alpha \Theta(x', y) \phi(x', y) \, dy$ and notice that $\chi \in L^1(\Omega)$. In fact,

$$\begin{aligned} \|\chi\|_{L^1(\Omega)} &= \int_\Omega \left| \int_0^\infty y^\alpha \Theta(x', y) \phi(x', y) \, dy \right| \, dx' \\ &\leq \int_{\mathcal{C}} |y^\alpha \Theta(x', y) \phi(x', y)| \, dx \leq \|\Theta\|_{L^2(y^\alpha, \mathcal{C})} \|\phi\|_{L^2(y^\alpha, \mathcal{C})} < \infty, \end{aligned}$$

where we have used that $\Theta \in \mathring{H}^1(y^\alpha, \mathcal{C})$. This, together with (3.5), yields

$$\begin{aligned} \int_{\mathcal{C}} y^\alpha q_{n_k} \Theta \phi \, dx &= \int_{\Omega} q_{n_k} \left(\int_0^\infty y^\alpha \Theta(x', y) \phi(x', y) \, dy \right) \, dx' = \int_{\Omega} q_{n_k} \chi \, dx' \\ &\rightarrow \int_{\Omega} q_{\delta, \rho} \chi \, dx' = \int_{\mathcal{C}} y^\alpha q_{\delta, \rho} \Theta \phi \, dx \end{aligned} \quad (3.8)$$

as $k \uparrow \infty$. As a consequence, when $k \uparrow \infty$, the term $\text{I}_k \rightarrow 0$.

Applying directly (3.6), we conclude, as $k \uparrow \infty$, that

$$\text{II}_k = \int_{\mathcal{C}} y^\alpha (\mathbf{A} \nabla E_{n_k} \cdot \nabla \phi + q_{\delta, \rho} \Theta \phi) \, dx \rightarrow \int_{\mathcal{C}} y^\alpha (\mathbf{A} \nabla \Theta \cdot \nabla \phi + q_{\delta, \rho} \Theta \phi) \, dx. \quad (3.9)$$

To control the term III_k , we proceed as follows. First, notice that

$$|\text{III}_k| = \left| \int_0^\infty y^\alpha \int_{\Omega} q_{n_k}(x') [E_{n_k}(x', y) - \Theta(x', y)] \phi(x', y) \, dx' \, dy \right| \leq \bar{q} \int_0^\infty \zeta_{n_k}(y) \, dy,$$

where

$$\zeta_{n_k}(y) := y^\alpha \|E_{n_k}(\cdot, y) - \Theta(\cdot, y)\|_{L^2(\Omega)} \|\phi(\cdot, y)\|_{L^2(\Omega)}.$$

Since $E_{n_k} \in \mathring{H}^1(y^\alpha, \mathcal{C})$, (2.9) implies that $\{E_{n_k}(\cdot, y)\} \subset H^{1/2}(\Omega)$ for a.e. $y \in (0, \infty)$. This and the trace estimate (2.11) allow us to conclude that $\{E_{n_k}(\cdot, y)\}$ converges to $\Theta(\cdot, y)$ in $L^2(\Omega)$ and thus that, a.e. $y \in (0, \infty)$, $\zeta_{n_k}(y) \rightarrow 0$ as $n_k \uparrow \infty$. In addition, it can be proved that $(\zeta_{n_k})_{k \in \mathbb{N}}$ is uniformly integrable and that, for every $\epsilon > 0$, there exists a set $A \subset (0, \infty)$ of finite measure such that, for all n_k , $\int_{A^c} \zeta_{n_k} \, dy < \epsilon$; the latter being a consequence of the exponential decay of E_{n_k} in the extended variable y [36, Proposition 3.1]. We can thus apply the Vitali convergence theorem [23, Section 6] to conclude that

$$\left| \int_0^\infty y^\alpha \int_{\Omega} q_{n_k}(x') [E_{n_k}(x', y) - \Theta(x', y)] \phi(x', y) \, dx' \, dy \right| \rightarrow 0, \quad (3.10)$$

as $k \uparrow \infty$.

It thus follows from (3.8)–(3.10) that

$$\int_{\mathcal{C}} y^\alpha (\mathbf{A} \nabla \Theta \cdot \nabla \phi + q_{\delta, \rho} \Theta \phi) \, dx = d_s \langle f, \text{tr}_\Omega \phi \rangle \quad \forall \phi \in \mathring{H}^1(y^\alpha, \mathcal{C}),$$

i.e., we have thus proved that $\Theta = E(q_{\delta, \rho})$.

We now invoke the fact that $\text{tr}_\Omega E_{n_k} \rightarrow \text{tr}_\Omega E(q_{\delta, \rho})$ in $L^2(\Omega)$ and that $\|\cdot\|_{L^2(\Omega)}$ is weakly lower semicontinuous to conclude that

$$\begin{aligned} \mathcal{J}_{\rho, \delta}(q_{\rho, \delta}) &= \frac{1}{2} \|\text{tr}_\Omega E(q_{\delta, \rho}) - z_\delta\|_{L^2(\Omega)}^2 + \frac{\rho}{2} \|q_{\delta, \rho} - q^*\|_{L^2(\Omega)}^2 \\ &\leq \liminf_{k \rightarrow \infty} \left(\frac{1}{2} \|\text{tr}_\Omega E_{n_k} - z_\delta\|_{L^2(\Omega)}^2 + \frac{\rho}{2} \|q_{n_k} - q^*\|_{L^2(\Omega)}^2 \right) \\ &= \inf_{q \in \mathcal{Q}} \left(\frac{1}{2} \|\text{tr}_\Omega E(q) - z_\delta\|_{L^2(\Omega)}^2 + \frac{\rho}{2} \|q - q^*\|_{L^2(\Omega)}^2 \right). \end{aligned}$$

Consequently, $q_{\delta, \rho}$ is a minimizer for problem (3.3). This concludes the proof. \square

REMARK 3.3 (non-uniqueness). Due to the lack of strict convexity of the cost functional $\mathcal{J}_{\delta, \rho}$, the minimizer $q_{\delta, \rho}$ may not be unique.

The previous remark motivates the following definition [44, Section 4.4].

DEFINITION 3.4 (local solution). A coefficient \tilde{q} is said to be a local solution for problem (3.3) if there exists $\epsilon > 0$ such that for all $q \in \mathcal{Q}$ that satisfies $\|q - \tilde{q}\|_{L^2(\Omega)} < \epsilon$ we have that $\mathcal{J}_{\rho,\delta}(\tilde{q}) \leq \mathcal{J}_{\rho,\delta}(q)$.

REMARK 3.5 (local solution). With the previous definition at hand, the results of Theorem 3.2 immediately yield the existence of a local solution $q_{\delta,\rho}$ for (3.3).

THEOREM 3.6 (existence of solutions). *The fractional regularization problem (1.6) has a solution $q_{\delta,\rho}$.*

Proof. The result follows directly from the results of Theorems 3.1 and 3.2. \square

3.2. Optimality conditions. We begin with a classical result.

LEMMA 3.7 (first-order optimality condition). *If $q_{\delta,\rho} \in \mathcal{Q}$ minimizes the functional $\mathcal{J}_{\delta,\rho}$, then $q_{\delta,\rho}$ solves the variational inequality*

$$(\mathcal{J}'_{\delta,\rho}(q_{\delta,\rho}), q - q_{\delta,\rho})_{L^2(\Omega)} \geq 0 \quad (3.11)$$

for every $q \in \mathcal{Q}$.

To explore (3.11), in what follows, we derive the Fréchet and thus the Gâteaux differentiability of the map E . To accomplish this task, we define, for $q \in \mathcal{Q}$, the map

$$\Psi : L^\infty(\Omega) \rightarrow \mathring{H}^1(y^\alpha, \mathcal{C}), \quad h \mapsto \Psi(h) = \psi, \quad (3.12)$$

where $\psi \in \mathring{H}^1(y^\alpha, \mathcal{C})$ solves

$$a_{\mathcal{C}}(\psi, \phi)(q) = - \int_{\mathcal{C}} y^\alpha h E(q) \phi \, dx \quad \forall \phi \in \mathring{H}^1(y^\alpha, \mathcal{C}). \quad (3.13)$$

Since $h \in L^\infty(\Omega)$ and $E(q) \in \mathring{H}^1(y^\alpha, \mathcal{C})$, a simple application of the Lax–Milgram Lemma reveals that there exist a unique ψ that solves (3.13) together with the estimate

$$\|\nabla \psi\|_{L^2(y^\alpha, \mathcal{C})} \lesssim \|h\|_{L^\infty(\Omega)} \|f\|_{\mathbb{H}^{-s}(\Omega)}. \quad (3.14)$$

Consequently, the map $\Psi : L^\infty(\Omega) \rightarrow \mathring{H}^1(y^\alpha, \mathcal{C})$ is linear and continuous.

LEMMA 3.8 (first-order Fréchet differentiability). Let $E : \mathcal{Q} \rightarrow \mathring{H}^1(y^\alpha, \mathcal{C})$ be the extended coefficient-to-solution operator defined in (3.1). The map E is first-order Fréchet differentiable. In addition, for $q \in \mathcal{Q}$ and $h \in L^\infty(\Omega)$, we have that $E'(q)h = \Psi(h)$, where Ψ is defined as in (3.12)–(3.13), and that

$$\|\nabla E'(q)h\|_{L^2(y^\alpha, \mathcal{C})} \lesssim \|f\|_{\mathbb{H}^{-s}(\Omega)} \|h\|_{L^\infty(\Omega)}, \quad (3.15)$$

where the hidden constant is independent of q and h .

Proof. Let $q \in \mathcal{Q}$ and consider $h \in L^\infty(\Omega)$ such that $q + h \in \mathcal{Q}$. In view of (2.14) and the definition of E , given by (3.1), we arrive at the identity

$$a_{\mathcal{C}}(E(q), \phi)(q) = d_s \langle f, \text{tr}_\Omega \phi \rangle = a_{\mathcal{C}}(E(q+h), \phi)(q+h) \quad \forall \phi \in \mathring{H}^1(y^\alpha, \mathcal{C}). \quad (3.16)$$

This, and the fact that ψ solves (3.13) allow us to conclude that

$$\begin{aligned} & \int_{\mathcal{C}} y^\alpha [\mathbf{A} \nabla (E(q+h) - E(q)) \cdot \nabla \phi + (q+h)(E(q+h) - E(q)) \phi] \, dx \\ &= - \int_{\mathcal{C}} y^\alpha h E(q) \phi \, dx = a_{\mathcal{C}}(\psi, \phi)(q) \quad \forall \phi \in \mathring{H}^1(y^\alpha, \mathcal{C}). \end{aligned} \quad (3.17)$$

Consequently, we can thus arrive at

$$a_{\mathcal{C}}(E(q+h) - E(q) - \Psi(h), \phi)(q+h) = - \int_{\mathcal{C}} y^\alpha h \Psi(h) \phi \, dx \quad \forall \phi \in \mathring{H}^1(y^\alpha, \mathcal{C}),$$

which implies that

$$\begin{aligned} \|\nabla(E(q+h) - E(q) - \Psi(h))\|_{L^2(y^\alpha, \mathcal{C})} &\lesssim \|h\|_{L^\infty(\Omega)} \|\Psi(h)\|_{L^2(y^\alpha, \mathcal{C})} \\ &\lesssim \|h\|_{L^\infty(\Omega)}^2 \|f\|_{\mathbb{H}^{-s}(\Omega)}, \end{aligned}$$

where, to obtain the last estimate, we have used (3.14).

Since $\Psi : L^\infty(\Omega) \rightarrow \mathring{H}^1(y^\alpha, \mathcal{C})$ is a linear and continuous map, we have thus obtained that the map E is first-order Fréchet differentiable and that $E'(q)h = \Psi(h)$. The estimate (3.15) follows from (3.14). This concludes the proof. \square

To describe the variational inequality (3.11), we introduce the adjoint variable

$$\mathcal{P} \in \mathring{H}^1(y^\alpha, \mathcal{C}) : \quad a_{\mathcal{C}}(w, \mathcal{P})(q) = d_s \langle \text{tr}_\Omega E(q) - z_\delta, w \rangle \quad \forall w \in \mathring{H}^1(y^\alpha, \mathcal{C}). \quad (3.18)$$

With this adjoint state at hand, we define the auxiliary variable

$$\mathfrak{p}(q) = -\frac{1}{d_s} \int_0^\infty y^\alpha E(q) \mathcal{P} \, dy. \quad (3.19)$$

The optimality conditions (3.11) in this setting now read as follow.

THEOREM 3.9 (first-order optimality condition). *If $q_{\delta, \rho} \in \mathcal{Q}$ minimizes the functional $\mathcal{J}_{\delta, \rho}$, then $q_{\delta, \rho}$ solves the variational inequality*

$$(\mathfrak{p}(q_{\delta, \rho}) + \rho(q_{\delta, \rho} - q^*), q - q_{\delta, \rho})_{L^2(\Omega)} \geq 0 \quad \forall q \in \mathcal{Q}, \quad (3.20)$$

where $\mathfrak{p}(q_{\delta, \rho}) \in L^2(\Omega)$ is defined in (3.19).

Proof. Since E is differentiable, an application of (3.11) reveals that

$$(\text{tr}_\Omega E(q_{\delta, \rho}) - z_\delta, \text{tr}_\Omega E'(q_{\delta, \rho})(q - q_{\delta, \rho}))_{L^2(\Omega)} + \rho(q_{\delta, \rho} - q^*, q - q_{\delta, \rho})_{L^2(\Omega)} \geq 0.$$

Invoking (3.13) and the results of Lemma 3.8, we obtain that $E'(q_{\delta, \rho})(q - q_{\delta, \rho})$ solves

$$a_{\mathcal{C}}(E'(q_{\delta, \rho})(q - q_{\delta, \rho}), w)(q_{\delta, \rho}) = - \int_{\mathcal{C}} y^\alpha (q - q_{\delta, \rho}) E(q_{\delta, \rho}) w \, dx \quad \forall w \in \mathring{H}^1(y^\alpha, \mathcal{C}). \quad (3.21)$$

By setting $w = E'(q_{\delta, \rho})(q - q_{\delta, \rho}) \in \mathring{H}^1(y^\alpha, \mathcal{C})$ in (3.18) and using first (3.21) and then (3.19) we arrive at

$$\begin{aligned} d_s \langle \text{tr}_\Omega E(q_{\delta, \rho}) - z_\delta, \text{tr}_\Omega E'(q_{\delta, \rho})(q - q_{\delta, \rho}) \rangle &= a_{\mathcal{C}}(E'(q_{\delta, \rho})(q - q_{\delta, \rho}), \mathcal{P}(q_{\delta, \rho}))(q_{\delta, \rho}) \\ &= - \int_{\mathcal{C}} y^\alpha (q - q_{\delta, \rho}) E(q_{\delta, \rho}) \mathcal{P}(q_{\delta, \rho}) \, dx = (d_s \mathfrak{p}(q_{\delta, \rho}), q - q_{\delta, \rho})_{L^2(\Omega)}. \end{aligned}$$

This concludes the proof. \square

In what follows we present higher-order differentiability results for the extended coefficient-to-solution operator E .

LEMMA 3.10 (high-order Fréchet differentiability). *The map $E : \mathcal{Q} \rightarrow \mathring{H}^1(y^\alpha, \mathcal{C})$ is infinitely Fréchet differentiable on the set $\mathcal{Q} \subset L^\infty(\Omega)$. In addition, for $q \in \mathcal{Q}$ and $m \geq 2$, the action of the m -th Fréchet derivative at q , $E^{(m)}(q)$, in the direction*

$(h_1, h_2, \dots, h_m) \in L^\infty(\Omega)^m$, that is denoted by $\psi = E^{(m)}(q)(h_1, h_2, \dots, h_m)$, corresponds to the unique solution to

$$\psi \in \hat{H}^1(y^\alpha, \mathcal{C}) : \quad a_{\mathcal{C}}(\psi, \phi) = - \sum_{i=1}^m \int_{\mathcal{C}} y^\alpha h_i \left[E^{(m-1)}(q) \xi_i \right] \phi \, dx \quad (3.22)$$

for all $\phi \in \hat{H}^1(y^\alpha, \mathcal{C})$, where $\xi_i := (h_1, \dots, h_{i-1}, h_{i+1}, \dots, h_m)$. Furthermore,

$$\|\nabla \psi\|_{L^2(y^\alpha, \mathcal{C})} \lesssim \|f\|_{\mathbb{H}^{-s}(\Omega)} \prod_{i=1}^m \|h_i\|_{L^\infty(\Omega)}, \quad (3.23)$$

where the hidden constant is independent of q and h .

Proof. On the basis of an induction argument, the proof follows the arguments developed in the proof of Lemma 3.8. For brevity we skip details. \square

In order to derive error estimates, in what follows we assume the following second-order optimality condition: If $q_{\delta, \rho}$ solves the variational inequality (3.11), then there exists $\theta > 0$ such that

$$\mathcal{J}''_{\delta, \rho}(q_{\delta, \rho})(q, q) \geq \theta \|q\|_{L^2(\Omega)}^2 \quad \forall q \in L^\infty(\Omega). \quad (3.24)$$

The following result guarantees that in a neighborhood of a local solution the second order derivatives of the functional $\mathcal{J}_{\delta, \rho}$ are coercive.

LEMMA 3.11 (local coercivity of $\mathcal{J}''_{\delta, \rho}$). *If $q_{\delta, \rho}$ denotes a local solution for problem (3.3), then there exists $\epsilon > 0$ such that for every \hat{q} in the neighborhood $\|\hat{q} - q_{\delta, \rho}\|_{L^2(\Omega)} \leq \epsilon$, we have that*

$$\mathcal{J}''_{\delta, \rho}(\hat{q})(q, q) \geq \frac{\theta}{2} \|q\|_{L^2(\Omega)}^2, \quad (3.25)$$

for all $q \in L^\infty(\Omega)$.

Proof. Notice that, in view of [31, Proposition 2.22] we have, for $q \in L^\infty(\Omega)$ and ϵ sufficiently small, that

$$\begin{aligned} \mathcal{J}''_{\delta, \rho}(\hat{q})(q, q) &= \mathcal{J}''_{\delta, \rho}(q_{\delta, \rho})(q, q) + [\mathcal{J}''_{\delta, \rho}(\hat{q})(q, q) - \mathcal{J}''_{\delta, \rho}(q_{\delta, \rho})(q, q)] \\ &\geq \theta \|q\|_{L^2(\Omega)}^2 - C\epsilon \|q\|_{L^2(\Omega)}^2 \geq \frac{\theta}{2} \|q\|_{L^2(\Omega)}^2, \end{aligned}$$

which yields the desired estimate (3.25). \square

3.3. A regularity result. Define

$$\Pi_{[0, \bar{q}]}(q)(x') := \min \{\bar{q}, \max \{0, q(x')\}\} \quad \text{for all } x' \text{ in } \bar{\Omega}. \quad (3.26)$$

In view of (3.20), standard argument allow us to conclude that

$$q_{\delta, \rho} = \Pi_{[0, \bar{q}]} \left(q^* - \frac{1}{\rho} \mathbf{p}(q_{\delta, \rho}) \right). \quad (3.27)$$

We present the following regularity result.

THEOREM 3.12 (regularity estimate). *Let $q_{\delta, \rho}$ be a local solution for (3.3). If $q^* \in H^1(\Omega)$, then $q_{\delta, \rho} \in H^1(\Omega)$. In addition, we have that*

$$\begin{aligned} \|\nabla_{x'} q_{\delta, \rho}\|_{L^2(\Omega)} + \|\nabla_{x'} \mathbf{p}(q_{\delta, \rho})\|_{L^2(\Omega)} &\lesssim \|\nabla_{x'} q^*\|_{L^2(\Omega)} \\ &\quad + \|f\|_{\mathbb{H}^{-s}(\Omega)} (\|f\|_{\mathbb{H}^{-s}(\Omega)} + \|z_\delta\|_{\mathbb{H}^{-s}(\Omega)}), \end{aligned} \quad (3.28)$$

where the hidden constant is independent of $q_{\delta,\rho}$ and the problem data.

Proof. Notice that, since $E(q_{\delta,\rho})$ and $\mathcal{P}(q_{\delta,\rho})$ belong to $\mathring{H}^1(y^\alpha, \mathcal{C})$, definition (3.19) implies that

$$\nabla_{x'} \mathbf{p}(q_{\delta,\rho}) = -\frac{1}{d_s} \int_0^\infty y^\alpha (\nabla_{x'} E(q_{\delta,\rho}) \mathcal{P}(q_{\delta,\rho}) + E(q_{\delta,\rho}) \nabla_{x'} \mathcal{P}(q_{\delta,\rho})) \, dy. \quad (3.29)$$

Stability estimates for the problems that $E(q_{\delta,\rho})$ and $\mathcal{P}(q_{\delta,\rho})$ solve reveal that

$$\begin{aligned} \|\nabla_{x'} \mathbf{p}(q_{\delta,\rho})\|_{L^2(\Omega)} &\lesssim \|\nabla E(q_{\delta,\rho})\|_{L^2(y^\alpha, \mathcal{C})} \|\nabla \mathcal{P}(q_{\delta,\rho})\|_{L^2(y^\alpha, \mathcal{C})} \\ &\lesssim \|f\|_{\mathbb{H}^{-s}(\Omega)} (\|f\|_{\mathbb{H}^{-s}(\Omega)} + \|z_\delta\|_{\mathbb{H}^{-s}(\Omega)}). \end{aligned}$$

The desired result thus follows from an application of [30, Theorem A.1]. \square

4. The truncated identification problem. Since the extended identification problem of section 3 is posed on the semi-infinite cylinder $\mathcal{C} = \Omega \times (0, \infty)$, a first step towards discretization consists in the truncation of the cylinder \mathcal{C} to the bounded domain $\mathcal{C}_\mathcal{Y} = \Omega \times (0, \mathcal{Y})$ and study the effect of truncation.

We begin our analysis by defining the weighted Sobolev space

$$\mathring{H}^1(y^\alpha, \mathcal{C}_\mathcal{Y}) = \{w \in H^1(y^\alpha, \mathcal{C}_\mathcal{Y}) : w = 0 \text{ on } \partial_L \mathcal{C}_\mathcal{Y} \cup \Omega_\mathcal{Y}\},$$

where $\Omega_\mathcal{Y} := \Omega \times \{\mathcal{Y}\}$. We define the bilinear form $a_\mathcal{Y} : \mathring{H}^1(y^\alpha, \mathcal{C}_\mathcal{Y}) \times \mathring{H}^1(y^\alpha, \mathcal{C}_\mathcal{Y})$ by

$$a_\mathcal{Y}(w, \phi)(q) = \int_{\mathcal{C}_\mathcal{Y}} y^\alpha (\mathbf{A} \nabla w \cdot \nabla \phi + qw\phi) \, dx.$$

With this setting at hand, we introduce the coefficient-to-solution operator:

$$H : \mathcal{Q} \rightarrow \mathring{H}^1(y^\alpha, \mathcal{C}_\mathcal{Y}) \quad (4.1)$$

which, given a coefficient q , associates to it the unique solution $v =: H(q)$ of problem

$$v \in \mathring{H}^1(y^\alpha, \mathcal{C}_\mathcal{Y}) : \quad a_\mathcal{Y}(v, \phi)(q) = d_s \langle f, \text{tr}_\Omega \phi \rangle \quad \forall \phi \in \mathring{H}^1(y^\alpha, \mathcal{C}_\mathcal{Y}). \quad (4.2)$$

Finally, we introduce the cost functional

$$\mathcal{R}_{\delta,\rho}(q) := \frac{1}{2} \|\text{tr}_\Omega H(q) - z_\delta\|_{L^2(\Omega)}^2 + \frac{\rho}{2} \|q - q^*\|_{L^2(\Omega)}^2. \quad (4.3)$$

The *truncated identification problem* thus reads as follows:

$$\min_{q \in \mathcal{Q}} \mathcal{R}_{\delta,\rho}(q). \quad (4.4)$$

To describe first-order optimality conditions we introduce the adjoint variable

$$p \in \mathring{H}^1(y^\alpha, \mathcal{C}_\mathcal{Y}) : \quad a_\mathcal{Y}(w, p)(q) = d_s \langle \text{tr}_\Omega H(q) - z_\delta, w \rangle \quad \forall w \in \mathring{H}^1(y^\alpha, \mathcal{C}_\mathcal{Y}), \quad (4.5)$$

and define

$$\mathbf{r}(q) = -\frac{1}{d_s} \int_0^\infty y^\alpha H(q) p \, dy. \quad (4.6)$$

The arguments elaborated in the proof of Theorems 3.2 and 3.9 allow us to obtain the following results.

THEOREM 4.1 (existence and first-order optimality condition). *The regularization problem (4.4) has a solution $r_{\delta,\rho}$. In addition, if $r_{\delta,\rho}$ minimizes (4.4), then $r_{\delta,\rho}$ solves the variational inequality*

$$(\mathfrak{r}(r_{\delta,\rho}) + \rho(r_{\delta,\rho} - q^*), q - r_{\delta,\rho})_{L^2(\Omega)} \geq 0 \quad \forall q \in \mathcal{Q}, \quad (4.7)$$

where $\mathfrak{r}(r_{\delta,\rho}) \in L^2(\Omega)$ is defined in (4.6).

REMARK 4.2 (local solution). In view of the results of Theorem 4.1 we conclude that problem (4.4) has a local solution $r_{\delta,\rho}$ in the sense of Definition 3.4.

The arguments elaborated in the proof of Theorem 3.12 allow us to immediately arrive at the following regularity estimate.

THEOREM 4.3 (regularity estimate). *Let $r_{\delta,\rho}$ be a local solution for (4.4). If $q^* \in H^1(\Omega)$, then $r_{\delta,\rho} \in H^1(\Omega)$. In addition, we have that*

$$\begin{aligned} \|\nabla_{x'} r_{\delta,\rho}\|_{L^2(\Omega)} + \|\nabla_{x'} \mathfrak{r}(r_{\delta,\rho})\|_{L^2(\Omega)} &\lesssim \|\nabla_{x'} q^*\|_{L^2(\Omega)} \\ &+ \|f\|_{\mathbb{H}^{-s}(\Omega)} (\|f\|_{\mathbb{H}^{-s}(\Omega)} + \|z_\delta\|_{\mathbb{H}^{-s}(\Omega)}), \end{aligned} \quad (4.8)$$

where the hidden constant is independent of $r_{\delta,\rho}$ and the problem data.

4.1. Auxiliary estimates. The following error estimates are instrumental for the error analysis that we will perform in section 4.3.

LEMMA 4.4 (exponential error estimate I). *Let $q \in \mathcal{Q}$ and $\gamma \geq 1$. If $\mathcal{U}(q)$ and $v(q)$ denote the solutions to problems (2.14) and (4.2), respectively, then*

$$\|\mathrm{tr}_\Omega(\mathcal{U}(q) - v(q))\|_{\mathbb{H}^s(\Omega)} \lesssim \|\nabla(\mathcal{U}(q) - v(q))\|_{L^2(y^\alpha, \mathcal{C})} \lesssim e^{-\sqrt{\lambda_1}\gamma/4} \|f\|_{\mathbb{H}^{-s}(\Omega)}, \quad (4.9)$$

where λ_1 corresponds to the first eigenvalue of operator \mathcal{L} .

Proof. We invoke the problems that $\mathcal{U}(q)$ and $v(q)$ solve to arrive at

$$\int_{\mathcal{C}_\gamma} y^\alpha (\mathbf{A} \nabla(\mathcal{U}(q) - v(q)) \cdot \nabla \phi + q(\mathcal{U}(q) - v(q))\phi) \, dx = 0 \quad \forall \phi \in \mathring{H}^1(y^\alpha, \mathcal{C}_\gamma).$$

Notice that we have used the trivial fact that $\phi \in \mathring{H}^1(y^\alpha, \mathcal{C}_\gamma)$ can be extended by zero to \mathcal{C} , which guarantees that $\phi \in \mathring{H}^1(y^\alpha, \mathcal{C})$. Consequently, we obtain that

$$\begin{aligned} \|\nabla(\mathcal{U}(q) - v(q))\|_{L^2(y^\alpha, \mathcal{C}_\gamma)}^2 &\lesssim a_\gamma(\mathcal{U}(q) - v(q), \mathcal{U}(q) - v(q)) \\ &= a_\gamma(\mathcal{U}(q) - v(q), \mathcal{U}(q) - \mathcal{E}) \quad \forall \mathcal{E} \in \mathring{H}^1(y^\alpha, \mathcal{C}_\gamma). \end{aligned}$$

This immediately yields the quasi-best approximation property

$$\|\nabla(\mathcal{U}(q) - v(q))\|_{L^2(y^\alpha, \mathcal{C}_\gamma)} \lesssim \inf\{\|\nabla(\mathcal{U}(q) - \mathcal{E})\|_{L^2(y^\alpha, \mathcal{C}_\gamma)}, \mathcal{E} \in \mathring{H}^1(y^\alpha, \mathcal{C}_\gamma)\}.$$

The right-hand side of the previous expression is bounded as in [36, Lemma 3.3]. In fact, it provides the estimate

$$\|\nabla(\mathcal{U}(q) - v(q))\|_{L^2(y^\alpha, \mathcal{C}_\gamma)} \lesssim e^{-\sqrt{\lambda_1}\gamma/4} \|f\|_{\mathbb{H}^{-s}(\Omega)},$$

which, combined with the fact that [36, Proposition 3.1]

$$\|\nabla \mathcal{U}(q)\|_{L^2(y^\alpha, \mathcal{C} \setminus \mathcal{C}_\gamma)} \lesssim e^{-\sqrt{\lambda_1}\gamma/2} \|f\|_{\mathbb{H}^{-s}(\Omega)},$$

allows us to conclude. \square

LEMMA 4.5 (exponential error estimate II). *Let $q \in \mathcal{Q}$ and $r \in L^\infty(\Omega)$. We thus have the following estimate:*

$$|\mathcal{J}'_{\delta,\rho}(q)r - \mathcal{R}'_{\delta,\rho}(q)r| \lesssim e^{-\sqrt{\lambda_1}\gamma/4} \|f\|_{\mathbb{H}^{-s}(\Omega)} (\|f\|_{\mathbb{H}^{-s}(\Omega)} + \|z_\delta\|_{\mathbb{H}^{-s}(\Omega)}) \|r\|_{L^2(\Omega)}.$$

The functionals $\mathcal{J}_{\delta,\rho}$ and $\mathcal{R}_{\delta,\rho}$ are defined by (3.2) and (4.3), respectively.

Proof. First, notice that, in view of definitions (3.19) and (4.6), we have that

$$\mathcal{J}'_{\delta,\rho}(q)r - \mathcal{R}'_{\delta,\rho}(q)r = (\mathbf{p}(q) - \mathbf{r}(q), r)_{L^2(\Omega)}.$$

It thus suffices to bound the term $\|\mathbf{p}(q) - \mathbf{r}(q)\|_{L^2(\Omega)}$. In fact, we have that

$$\begin{aligned} \|\mathbf{p}(q) - \mathbf{r}(q)\|_{L^2(\Omega)} &= \frac{1}{d_s} \left\| \int_0^\infty y^\alpha [E(q)\mathcal{P}(q) - H(q)p(q)] \, dy \right\|_{L^2(\Omega)} \\ &= \frac{1}{d_s} \left\| \int_0^\infty y^\alpha [(E(q) - H(q))\mathcal{P}(q) - H(q)(p(q) - \mathcal{P}(q))] \, dy \right\|_{L^2(\Omega)}. \end{aligned}$$

Standard estimates allow us to arrive at

$$\begin{aligned} \|\mathbf{p}(q) - \mathbf{r}(q)\|_{L^2(\Omega)} &\lesssim \|\mathcal{P}(q)\|_{L^2(y^\alpha, \mathcal{C})} \|E(q) - H(q)\|_{L^2(y^\alpha, \mathcal{C})} \\ &\quad + \|H(q)\|_{L^2(y^\alpha, \mathcal{C})} \|p(q) - \mathcal{P}(q)\|_{L^2(y^\alpha, \mathcal{C})}. \end{aligned}$$

This, in view of the Poincaré inequality (2.10), the exponential error estimate (4.9), and stability estimates for the problems that $E(q) - H(q)$ and $p(q) - \mathcal{P}(q)$ solve allow us to conclude that

$$\begin{aligned} \|\mathbf{p}(q) - \mathbf{r}(q)\|_{L^2(\Omega)} &\lesssim e^{-\sqrt{\lambda_1}\gamma/4} \operatorname{tr}_\Omega E(q) - z_\delta \|_{\mathbb{H}^{-s}(\Omega)} \|f\|_{\mathbb{H}^{-s}(\Omega)} \\ &\quad + \|f\|_{\mathbb{H}^{-s}(\Omega)} \|\operatorname{tr}_\Omega [H(q) - E(q)]\|_{\mathbb{H}^{-s}(\Omega)}, \quad (4.10) \end{aligned}$$

where $(\lambda_1)^{\frac{1}{2}} = (\lambda_1(q))^{\frac{1}{2}}$ denotes the first eigenvalue of the operator \mathcal{L} . Applying (4.9), again, and stability estimates, we conclude that

$$\|\mathbf{p}(q) - \mathbf{r}(q)\|_{L^2(\Omega)} \lesssim e^{-\sqrt{\lambda_1}\gamma/4} \|f\|_{\mathbb{H}^{-s}(\Omega)} (\|f\|_{\mathbb{H}^{-s}(\Omega)} + \|z_\delta\|_{\mathbb{H}^{-s}(\Omega)}). \quad (4.11)$$

This implies the desired estimate and concludes the proof. \square

To state the following result, we define, for $q \in \mathcal{Q}$ and a.e. $x' \in \Omega$,

$$\mathbf{e}(q)(x') := \int_0^\infty y^\alpha E(q)(x', y)^2 \, dy. \quad (4.12)$$

Notice that, for every $q \in \mathcal{Q}$, $\mathbf{e}(q) \in L^1(\Omega)$ and $\|\mathbf{e}(q)\|_{L^1(\Omega)} \leq \|E(q)\|_{L^2(y^\alpha, \mathcal{C})}^2$. We present the following result.

LEMMA 4.6 (stability estimate). *Let $q, r \in \mathcal{Q}$ and E be the extended coefficient-to-solution operator defined in (3.1). If, for every $q \in \mathcal{Q}$, $\mathbf{e}(q) \in L^\infty(\Omega)$, then*

$$\|\operatorname{tr}_\Omega (\mathcal{U}(q) - \mathcal{U}(r))\|_{\mathbb{H}^s(\Omega)} \lesssim \|\nabla(\mathcal{U}(q) - \mathcal{U}(r))\|_{L^2(y^\alpha, \mathcal{C})} \lesssim \|\mathbf{e}(r)\|_{L^\infty(\Omega)}^{\frac{1}{2}} \|q - r\|_{L^2(\Omega)},$$

where $\mathcal{U}(q) = E(q)$ and $\mathcal{U}(r) = E(r)$ and the hidden constant is independent of E , q , r , and f .

Proof. Since $\mathcal{U}(q) - \mathcal{U}(r) \in \mathring{H}^1(y^\alpha, \mathcal{C})$, the first estimate follows immediately from the trace estimate (2.11). The remaining estimate follows upon exploiting the problems that $\mathcal{U}(q)$ and $\mathcal{U}(r)$ solve. In fact, notice that $\mathcal{U}(q) - \mathcal{U}(r)$ satisfies

$$\int_{\mathcal{C}} y^\alpha [\mathbf{A} \nabla(\mathcal{U}(q) - \mathcal{U}(r)) \cdot \nabla \phi + q(\mathcal{U}(q) - \mathcal{U}(r))\phi] dx + \int_{\mathcal{C}} y^\alpha (q - r)\mathcal{U}(r)\phi dx = 0$$

for all $\phi \in \mathring{H}^1(y^\alpha, \mathcal{C})$. We can thus set $\phi = \mathcal{U}(q) - \mathcal{U}(r) \in \mathring{H}^1(y^\alpha, \mathcal{C})$ and obtain, on the basis of (2.13), Fubini's theorem, and the Cauchy–Schwarz inequality that

$$\begin{aligned} & \|\nabla(\mathcal{U}(q) - \mathcal{U}(r))\|_{L^2(y^\alpha, \mathcal{C})}^2 \\ & \lesssim \|q - r\|_{L^2(\Omega)} \left\| \int_0^\infty y^\alpha |\mathcal{U}(r)(x', y)| |\mathcal{U}(q)(x', y) - \mathcal{U}(r)(x', y)| dy \right\|_{L^2(\Omega)} \\ & \leq \|q - r\|_{L^2(\Omega)} \left(\int_\Omega \mathbf{e}(r)(x') \left[\int_0^\infty y^\alpha |\mathcal{U}(q)(x', y) - \mathcal{U}(r)(x', y)|^2 dy \right] dx' \right)^{\frac{1}{2}}. \end{aligned}$$

Thus,

$$\|\nabla(\mathcal{U}(q) - \mathcal{U}(r))\|_{L^2(y^\alpha, \mathcal{C})}^2 \lesssim \|q - r\|_{L^2(\Omega)} \|\mathbf{e}(r)\|_{L^\infty(\Omega)}^{\frac{1}{2}} \|\mathcal{U}(q) - \mathcal{U}(r)\|_{L^2(y^\alpha, \mathcal{C})},$$

which in view of the Poincaré inequality (2.10) allows us to arrive at the desired estimate. \square

4.2. Convergence. The following result is important since guarantees that every local minimum of problem (3.3) can be approximated by local minima of (4.4).

THEOREM 4.7 (convergence result). *Let $q_{\delta, \rho} \in \mathcal{Q}$ be a local solution to (3.3). Then, there exists a sequence $\{r_{\delta, \rho}\}$ of local minima for (4.4) such that*

$$\{r_{\delta, \rho}\} \rightarrow q_{\delta, \rho} \tag{4.13}$$

as the truncation parameter $\mathcal{Y} \uparrow \infty$.

Proof. We begin by noticing that there exists $\epsilon > 0$ such that $q_{\delta, \rho}$ is the unique strict solution to the following problem:

$$\min_{q \in \mathcal{Q}(q_{\delta, \rho})} \mathcal{J}_{\delta, \rho}(q), \quad \mathcal{Q}(q_{\delta, \rho}) = \{q \in \mathcal{Q} : \|q - q_{\delta, \rho}\|_{L^2(\Omega)} \leq \epsilon\}. \tag{4.14}$$

Let us now consider the following truncated optimization problem over $\mathcal{Q}(q_{\delta, \rho})$:

$$\min_{r \in \mathcal{Q}(q_{\delta, \rho})} \mathcal{R}_{\delta, \rho}(r). \tag{4.15}$$

It is immediate that $\mathcal{Q}(q_{\delta, \rho}) \neq \emptyset$. This implies that (4.15) has at least one solution. Let $r_{\delta, \rho} := r_{\delta, \rho}(\mathcal{Y})$ be a solution to (4.15) for $\mathcal{Y} \geq \mathcal{Y}_0 > 1$. Notice that $\{r_{\delta, \rho}\}$ is a bounded sequence in $L^\infty(\Omega)$. As in the proof of Theorem 3.2 we deduce the existence of non-relabeled subsequence $\{r_{\delta, \rho}\}$ that converges weakly* to \tilde{q} in $L^\infty(\Omega)$. Thus

$$\begin{aligned} \mathcal{J}_{\delta, \rho}(q_{\delta, \rho}) & \leq \mathcal{J}_{\delta, \rho}(\tilde{q}) \leq \liminf_{\mathcal{Y} \rightarrow \infty} \mathcal{J}_{\delta, \rho}(r_{\delta, \rho}) = \liminf_{\mathcal{Y} \rightarrow \infty} \mathcal{R}_{\delta, \rho}(r_{\delta, \rho}) \leq \limsup_{\mathcal{Y} \rightarrow \infty} \mathcal{R}_{\delta, \rho}(r_{\delta, \rho}) \\ & \leq \limsup_{\mathcal{Y} \rightarrow \infty} \mathcal{R}_{\delta, \rho}(q_{\delta, \rho}) = \mathcal{J}_{\delta, \rho}(q_{\delta, \rho}). \end{aligned}$$

Since (4.14) has a unique solution, we can thus conclude that the sequence $\{r_{\delta, \rho}\}$ converges strongly in $L^2(\Omega)$ to the solution to (4.14). This concludes the proof. \square

4.3. Error estimates. The next results shows how a sequence of local minima of (4.4) $\{r_{\delta,\rho}\}$ approximates a local solution $q_{\delta,\rho}$ of problem (3.3).

THEOREM 4.8 (exponential error estimate). *Let $q_{\delta,\rho}$ be a local solution of problem (3.3). If $\{r_{\delta,\rho}\}$ denotes a sequence of local minima of (4.4) that converges to $q_{\delta,\rho}$ as $\mathcal{Y} \uparrow \infty$ in $L^2(\Omega)$, then*

$$\|q_{\delta,\rho} - r_{\delta,\rho}\|_{L^2(\Omega)} \lesssim e^{-\kappa\mathcal{Y}/4} \|f\|_{\mathbb{H}^{-s}(\Omega)} \left(\|f\|_{\mathbb{H}^{-s}(\Omega)} + \|z_\delta\|_{\mathbb{H}^{-s}(\Omega)} \right), \quad (4.16)$$

where $\kappa = (\lambda_1(r_{\delta,\rho}))^{\frac{1}{2}}$ and $\lambda_1(r_{\delta,\rho})$ denotes the first eigenvalue of the operator \mathcal{L} with q replaced by $r_{\delta,\rho}$. The hidden constant is independent of $q_{\delta,\rho}$, $r_{\delta,\rho}$, f , and z_δ .

Proof. In view of the result of Lemma 3.11, we choose $\epsilon > 0$ sufficiently small such that, for \hat{q} in the neighborhood $\|\hat{q} - q_{\delta,\rho}\|_{L^2(\Omega)} \leq \epsilon$, the following estimate holds:

$$\mathcal{J}_{\delta,\rho}''(\hat{q})(q, q) \geq \frac{\theta}{2} \|q\|_{L^2(\Omega)}^2 \quad \forall q \in L^\infty(\Omega). \quad (4.17)$$

Since the sequence $\{r_{\delta,\rho}\}$ converges to $q_{\delta,\rho}$ in $L^2(\Omega)$ as $\mathcal{Y} \uparrow \infty$, we deduce the existence of \mathcal{Y}_0 such that for $\mathcal{Y} \geq \mathcal{Y}_0$, we have that $\|q_{\delta,\rho} - r_{\delta,\rho}\|_{L^2(\Omega)} \leq \epsilon$. We are thus able to set, for $\mathcal{Y} \geq \mathcal{Y}_0$ and $\zeta \in [0, 1]$, $q = r_{\delta,\rho} - q_{\delta,\rho}$ and $\hat{q} = \zeta q_{\delta,\rho} + (1 - \zeta)r_{\delta,\rho}$ into the estimate (4.17) and obtain that

$$\begin{aligned} \frac{\theta}{2} \|r_{\delta,\rho} - q_{\delta,\rho}\|_{L^2(\Omega)}^2 &\leq \mathcal{J}_{\delta,\rho}''(\hat{q})(r_{\delta,\rho} - q_{\delta,\rho}, r_{\delta,\rho} - q_{\delta,\rho}) \\ &= \mathcal{J}_{\delta,\rho}'(r_{\delta,\rho})(r_{\delta,\rho} - q_{\delta,\rho}) - \mathcal{J}_{\delta,\rho}'(q_{\delta,\rho})(r_{\delta,\rho} - q_{\delta,\rho}). \end{aligned}$$

Notice that $\|\hat{q} - q_{\delta,\rho}\|_{L^2(\Omega)} = (1 - \zeta)\|q_{\delta,\rho} - r_{\delta,\rho}\|_{L^2(\Omega)} \leq \epsilon$.

In view of (3.11) we immediately conclude that $-\mathcal{J}_{\delta,\rho}'(q_{\delta,\rho})(r_{\delta,\rho} - q_{\delta,\rho}) \leq 0$. On the other hand, (4.7) reveals that $-\mathcal{R}'_{\delta,\rho}(r_{\delta,\rho})(r_{\delta,\rho} - q_{\delta,\rho}) \geq 0$. Consequently,

$$\frac{\theta}{2} \|r_{\delta,\rho} - q_{\delta,\rho}\|_{L^2(\Omega)}^2 \leq (\mathcal{J}'_{\delta,\rho}(r_{\delta,\rho}), r_{\delta,\rho} - q_{\delta,\rho})_{L^2(\Omega)} - (\mathcal{R}'_{\delta,\rho}(r_{\delta,\rho}), r_{\delta,\rho} - q_{\delta,\rho})_{L^2(\Omega)}.$$

We now exploit that $\mathcal{J}'_{\delta,\rho}(r_{\delta,\rho}) = \mathbf{p}(r_{\delta,\rho}) + \rho(r_{\delta,\rho} - q^*)$ and that $\mathcal{R}'_{\delta,\rho}(r_{\delta,\rho}) = \mathbf{r}(r_{\delta,\rho}) + \rho(r_{\delta,\rho} - q^*)$ to obtain the following estimate

$$\begin{aligned} \frac{\theta}{2} \|r_{\delta,\rho} - q_{\delta,\rho}\|_{L^2(\Omega)}^2 &\leq (\mathbf{p}(r_{\delta,\rho}) + \rho(r_{\delta,\rho} - q^*), r_{\delta,\rho} - q_{\delta,\rho})_{L^2(\Omega)} \\ &\quad - (\mathbf{r}(r_{\delta,\rho}) + \rho(r_{\delta,\rho} - q^*), r_{\delta,\rho} - q_{\delta,\rho})_{L^2(\Omega)} = (\mathbf{p}(r_{\delta,\rho}) - \mathbf{r}(r_{\delta,\rho}), r_{\delta,\rho} - q_{\delta,\rho})_{L^2(\Omega)}. \end{aligned}$$

The control of the term $(\mathbf{p}(r_{\delta,\rho}) - \mathbf{r}(r_{\delta,\rho}), r_{\delta,\rho} - q_{\delta,\rho})_{L^2(\Omega)}$ follows from Lemma 4.5. In fact, we have that

$$\|\mathbf{p}(r_{\delta,\rho}) - \mathbf{r}(r_{\delta,\rho})\|_{L^2(\Omega)} \lesssim e^{-\kappa\mathcal{Y}/4} \|f\|_{\mathbb{H}^{-s}(\Omega)} \left(\|f\|_{\mathbb{H}^{-s}(\Omega)} + \|z_\delta\|_{\mathbb{H}^{-s}(\Omega)} \right).$$

This concludes the proof. \square

THEOREM 4.9 (exponential error estimate). *Let $q_{\delta,\rho}$ be a local solution of problem (3.3). If $\{r_{\delta,\rho}\}$ denotes a sequence of local minima of (4.4) that converges to $q_{\delta,\rho}$ as $\mathcal{Y} \uparrow \infty$ in $L^2(\Omega)$, then*

$$\begin{aligned} \|\nabla(\mathcal{W}(q_{\delta,\rho}) - v(r_{\delta,\rho}))\|_{L^2(\mathcal{Y}^\alpha, \mathcal{C})} \\ \lesssim e^{-\kappa\mathcal{Y}/4} \|f\|_{\mathbb{H}^{-s}(\Omega)} \left(1 + \|\mathbf{e}(r_{\delta,\rho})\|_{L^\infty(\Omega)}^{\frac{1}{2}} \left(\|f\|_{\mathbb{H}^{-s}(\Omega)} + \|z_\delta\|_{\mathbb{H}^{-s}(\Omega)} \right) \right), \quad (4.18) \end{aligned}$$

where $\kappa = (\lambda_1(r_{\delta,\rho}))^{\frac{1}{2}}$ and $\lambda_1(r_{\delta,\rho})$ denotes the first eigenvalue of the operator \mathcal{L} with q replaced by $r_{\delta,\rho}$. The hidden constant is independent of $q_{\delta,\rho}$, $r_{\delta,\rho}$, f , and z_δ .

Proof. To derive the estimate (4.18), we proceed as follows:

$$\begin{aligned} \|\nabla(\mathcal{W}(q_{\delta,\rho}) - v(r_{\delta,\rho}))\|_{L^2(y^\alpha, \mathcal{C})} &\leq \|\nabla(\mathcal{W}(q_{\delta,\rho}) - \mathcal{W}(r_{\delta,\rho}))\|_{L^2(y^\alpha, \mathcal{C})} \\ &\quad + \|\nabla(\mathcal{W}(r_{\delta,\rho}) - v(r_{\delta,\rho}))\|_{L^2(y^\alpha, \mathcal{C})} = \text{I} + \text{II}. \end{aligned} \quad (4.19)$$

The control for the term I follows from applying, first, the estimate of Lemma 4.6 and then (4.16). In fact, we have that

$$\begin{aligned} \text{I} &\lesssim \|\mathbf{e}(r_{\delta,\rho})\|_{L^\infty(\Omega)}^{\frac{1}{2}} \|q_{\delta,\rho} - r_{\delta,\rho}\|_{L^2(\Omega)} \\ &\lesssim e^{-\kappa\mathcal{Y}/4} \|\mathbf{e}(r_{\delta,\rho})\|_{L^\infty(\Omega)}^{\frac{1}{2}} \|f\|_{\mathbb{H}^{-s}(\Omega)} (\|f\|_{\mathbb{H}^{-s}(\Omega)} + \|z_\delta\|_{\mathbb{H}^{-s}(\Omega)}). \end{aligned}$$

The estimate for the second term II follows from (4.9):

$$\text{II} \lesssim e^{-\kappa\mathcal{Y}/4} \|f\|_{\mathbb{H}^{-s}(\Omega)}.$$

A collection of the estimates for I and II yield the desired result. \square

5. A discretization scheme. In this section we present a fully discrete scheme that approximate solutions to the fractional identification problem (1.6). In view of the localization results of Theorem 3.1 and the exponential error estimates of Theorem 4.8 and 4.9, in what follows, we design and analyze an efficient solution technique to solve the truncated identification problem (4.4). We begin by introducing ingredients for a suitable finite element discretization of the truncated equation (4.2).

5.1. Finite element methods. We follow [2] and introduce a finite element technique that is based on the tensorization of a first-degree FEM in Ω with a suitable hp -FEM on the extended domain $(0, \mathcal{Y})$. The scheme achieves log-linear complexity with respect to the number of degrees of freedom in the domain Ω . To present it, we first introduce, on the extended interval $[0, \mathcal{Y}]$, the following geometric meshes with M elements and grading factor $\sigma \in (0, 1)$:

$$\mathcal{G}_\sigma^M = \{I_m\}_{m=1}^M, \quad I_1 = [0, \mathcal{Y}\sigma^{M-1}], \quad I_i = [\mathcal{Y}\sigma^{M-i+1}, \mathcal{Y}\sigma^{M-i}], \quad (5.1)$$

with $i \in \{2, \dots, M\}$. The main motivation for considering the meshes \mathcal{G}_σ^M , that are refined towards $y = 0$, is to compensate the rather singular behavior of \mathcal{W} , solution to problem (2.14), as $y \downarrow 0$; see [36, Theorem 2.7] and [2, Theorem 4.7]. On these meshes, we consider a *linear degree vector* $\mathbf{r} = (r_1, \dots, r_M) \in \mathbb{N}^M$ with slope \mathfrak{s} : $r_i := \max\{1, \lceil \mathfrak{s}i \rceil\}$, where $i = 1, 2, \dots, M$. We thus introduce the finite element space

$$S^{\mathbf{r}}((0, \mathcal{Y}), \mathcal{G}_\sigma^M) = \{v_M \in C[0, \mathcal{Y}] : v_M|_{I_m} \in \mathbb{P}_{r_m}(I_m), I_m \in \mathcal{G}_\sigma^M, m = 1, \dots, M\},$$

and the subspace of $S^{\mathbf{r}}((0, \mathcal{Y}), \mathcal{G}_\sigma^M)$ containing functions that vanish at $y = \mathcal{Y}$:

$$S_{\{\mathcal{Y}\}}^{\mathbf{r}}((0, \mathcal{Y}), \mathcal{G}_\sigma^M) = \{v_M \in S^{\mathbf{r}}((0, \mathcal{Y}), \mathcal{G}_\sigma^M) : v_M(\mathcal{Y}) = 0\}.$$

Let $\mathcal{T} = \{K\}$ be a conforming partition of $\bar{\Omega}$ into simplices K . We denote by \mathbb{T} a collection of conforming and shape regular meshes that are refinements of an original mesh \mathcal{T}_0 . For $\mathcal{T} \in \mathbb{T}$, we define $h_{\mathcal{T}} = \max\{\text{diam}(K) : K \in \mathcal{T}\}$ and $N = \#\mathcal{T}$, the number of degrees of freedom of \mathcal{T} . We introduce the finite element space

$$S_0^1(\Omega, \mathcal{T}) = \{v_h \in C(\bar{\Omega}) : v_h|_K \in \mathbb{P}_1(K) \quad \forall K \in \mathcal{T}, v_h|_{\partial\Omega} = 0\}.$$

With the meshes \mathcal{G}_σ^M and \mathcal{T} at hand, we define $\mathcal{T}_\gamma = \mathcal{T} \otimes \mathcal{G}_\sigma^M$ and the finite-dimensional *tensor product space*

$$\mathbb{V}_{N,M}^{1,\mathbf{r}}(\mathcal{T}_\gamma) := S_0^1(\Omega, \mathcal{T}) \otimes S_{\{\gamma\}}^{\mathbf{r}}((0, \gamma), \mathcal{G}_\sigma^M) \subset \mathring{H}^1(y^\alpha, \mathcal{C}). \quad (5.2)$$

We write $\mathbb{V}(\mathcal{T}_\gamma)$ if the arguments are clear from the context.

With this discrete setting at hand, we define the finite element approximation $V \in \mathbb{V}(\mathcal{T}_\gamma)$ of the solution $v \in \mathring{H}^1(y^\alpha, \mathcal{C}_\gamma)$ to problem (4.2) as follows:

$$V \in \mathbb{V}(\mathcal{T}_\gamma) : \quad a_{\mathcal{C}}(V, W)(q) = d_s \langle f, \text{tr}_\Omega W \rangle \quad \forall W \in \mathbb{V}(\mathcal{T}_\gamma). \quad (5.3)$$

The following a priori error estimate can be obtained [2, Theorem 5.4].

THEOREM 5.1 (a priori error estimate). *Let \mathcal{G}_σ^M be the geometric meshes defined in (5.1), where $\gamma \sim |\log h_{\mathcal{T}}|$ with a sufficiently large implied constant. If $V \in \mathbb{V}(\mathcal{T}_\gamma)$ denotes the solution to (5.3) then, there exists a minimal slope \mathfrak{s}_{\min} , independent of $h_{\mathcal{T}}$ and f , such that for linear degree vectors \mathbf{r} with slope $\mathfrak{s} \geq \mathfrak{s}_{\min}$ there holds*

$$\|u - \text{tr}_\Omega V\|_{\mathbb{H}^s(\Omega)} \lesssim \|\nabla(\mathcal{U} - V)\|_{L^2(y^\alpha, \mathcal{C})} \lesssim h_{\mathcal{T}} \|f\|_{\mathbb{H}^{1-s}(\Omega)}. \quad (5.4)$$

The hidden constant is independent of u , \mathcal{U} , V , f and the discretization parameters.

5.2. A fully discrete scheme. We begin by defining the discrete sets

$$\mathbb{Q}(\mathcal{T}) = \{Q \in L^\infty(\Omega) : Q|_K \in \mathbb{P}_0(K) \quad \forall K \in \mathcal{T}_\Omega\}, \quad \mathbb{Q}_{ad}(\mathcal{T}) = \mathbb{Q}(\mathcal{T}) \cap \mathcal{Q},$$

and the discrete coefficient-to-solution operator $F : \mathcal{Q} \rightarrow \mathbb{V}(\mathcal{T}_\gamma)$, which associates to an element $q \in \mathcal{Q}$ the unique discrete solution $V =: F(q)$ of problem (5.3). With this operator at hand, we define the discrete reduced cost functional

$$\mathcal{D}_{\delta, \rho}(Q) := \frac{1}{2} \|\text{tr}_\Omega F(Q) - z_\delta\|_{L^2(\Omega)}^2 + \frac{\rho}{2} \|Q - q^*\|_{L^2(\Omega)}^2, \quad (5.5)$$

and the following fully discrete approximation of the identification problem (4.4):

$$\min_{Q \in \mathbb{Q}_{ad}(\mathcal{T})} \mathcal{D}_{\delta, \rho}(Q). \quad (5.6)$$

LEMMA 5.2 (existence of discrete solutions). *The discrete problem (5.6) has a solution $Q_{\delta, \rho}$.*

Proof. Let $(Q_n)_{n \in \mathbb{N}} \subset \mathbb{Q}_{ad}(\mathcal{T})$ be a minimizing sequence for problem (5.6). We thus have the existence of a nonrelabeled subsequence of $(Q_n)_{n \in \mathbb{N}}$ that converges, in the $L^2(\Omega)$ -norm, to an element Q of the finite dimensional space $\mathbb{Q}_{ad}(\mathcal{T})$. Standard arguments reveal that the sequence $(F(Q_n))_{n \in \mathbb{N}}$ converges to $F(Q)$ in the space $\mathbb{V}(\mathcal{T}_\gamma)$. This concludes the proof. \square

In order to present the following result, we define the set

$$\mathcal{I}(u^\dagger) := \{q \in \mathcal{Q} \mid U(q) = u^\dagger\}, \quad (5.7)$$

In (5.7), U denotes the coefficient-to-solution operator associated to problem (1.3). Notice that, in view of Theorem 3.1, we have that $\mathcal{I}(u^\dagger) = \{q \in \mathcal{Q} \mid \text{tr}_\Omega E(q) = u^\dagger\}$.

We now introduce the concept of q^* -minimum-norm solution for the fractional identification problem (1.6).

LEMMA 5.3 (q^* -minimum-norm solution). *The optimization problem*

$$\min_{q \in \mathcal{I}(u^\dagger)} \|q - q^*\|_{L^2(\Omega)}^2 \quad (5.8)$$

attains a unique solution q^\dagger , which is called the q^* -minimum–norm solution of problem (1.6).

Proof. Notice that, in view of the arguments elaborated in the proof of Theorem 3.2, we can conclude that $\mathcal{I}(u^\dagger)$, defined in (5.7), is a nonempty set that is weakly closed in $L^2(\Omega)$. In addition, it is immediate that $\mathcal{I}(u^\dagger)$ is a convex and bounded subset of $L^2(\Omega)$. Consequently, $\mathcal{I}(u^\dagger)$ is weakly sequentially compact in $L^2(\Omega)$ [17, Theorem 5.50]. Since $\|\cdot\|_{L^2(\Omega)}$ is weakly lower semicontinuous, an application of [17, Theorem 5.51] allows us to conclude. \square

Since it will be instrumental in the analysis that we will perform, we introduce the $L^2(\Omega)$ –orthogonal projection operator $\pi_{x'} : L^2(\Omega) \rightarrow \mathbb{Q}(\mathcal{T})$ as follows: For $K \in \mathcal{T}$ and $q \in L^2(\Omega)$, $\pi_{x'}$ is defined as [21, Section 1.63]

$$\pi_{x'} q|_K = \frac{1}{|K|} \int_K q(x') \, dx'. \quad (5.9)$$

Notice that $\pi_{x'} \mathcal{Q} \subset \mathbb{Q}_{ad}(\mathcal{T}_\Omega)$. In addition, for $1 \leq p \leq \infty$, $s \in (0, 1]$, and $q \in W^{s,p}(\Omega)$, we have the following error estimate [21, Proposition 1.135]:

$$\|q - \pi_{x'} q\|_{L^p(\Omega)} \lesssim h_{\mathcal{T}}^s |q|_{W^{s,p}(\Omega)} \quad (5.10)$$

In the following result we show that the finite element solutions of problem (5.6) converge to q^* -minimum–norm solution q^\dagger .

THEOREM 5.4 (convergence of solutions). *Let $(\mathcal{T}_{h_n})_n$ be a sequence of conforming, shape–regular, and quasi–uniform meshes of Ω and let $h_n := h_{\mathcal{T}_n}$ be the meshwidth of \mathcal{T}_n . Assume that there exists $0 < \gamma < 1$ such that $q^\dagger \in W^{\gamma,\infty}(\Omega)$. Let $(\delta_n)_{n \in \mathbb{N}}$ be a sequence in \mathbb{R}^+ , and consider $\rho_n := \rho_{\delta_n, h_n}$ to be such that*

$$\rho_n \rightarrow 0, \quad \frac{\delta_n^2}{\rho_n} \rightarrow 0, \quad \text{and} \quad \frac{h_n^{2\gamma}}{\rho_n} \rightarrow 0 \quad \text{as } n \uparrow \infty. \quad (5.11)$$

Let $(z_n)_{n \in \mathbb{N}} = (z_{\delta_n})_{n \in \mathbb{N}}$ be a sequence in $L^2(\Omega)$ such that $\|u^\dagger - z_n\|_{L^2(\Omega)} \leq \delta_n$ and let Q_n be a minimizer of the following problem (cf. (5.6)):

$$\min_{Q \in \mathbb{Q}_{ad}(\mathcal{T}_\Omega)} \mathcal{D}_{\delta_n, \rho_n}(Q) := \frac{1}{2} \|\text{tr}_\Omega F(Q) - z_n\|_{L^2(\Omega)}^2 + \frac{\rho_n}{2} \|Q - q^*\|_{L^2(\Omega)}^2. \quad (5.12)$$

Then the sequence $(Q_n)_{n \in \mathbb{N}}$ converges to q^\dagger in $L^2(\Omega)$ as $n \uparrow \infty$.

Proof. Since Q_n is optimal for problem (5.12), we immediately arrive at

$$\mathcal{D}_{\delta_n, \rho_n}(Q_n) \leq \frac{1}{2} \|\text{tr}_\Omega F(\pi_{x'} q^\dagger) - z_n\|_{L^2(\Omega)}^2 + \frac{\rho_n}{2} \|\pi_{x'} q^\dagger - q^*\|_{L^2(\Omega)}^2, \quad (5.13)$$

where $\pi_{x'}$ is defined as in (5.9).

We now proceed to estimate the first term on the right–hand side of (5.13). In view of the estimates (5.4) and $\|u^\dagger - z_n\|_{L^2(\Omega)} \leq \delta_n$, we obtain that

$$\begin{aligned} \|\text{tr}_\Omega F(\pi_{x'} q^\dagger) - z_n\|_{L^2(\Omega)} &\leq \|u(\pi_{x'} q^\dagger) - \text{tr}_\Omega F(\pi_{x'} q^\dagger)\|_{L^2(\Omega)} + \|u(q^\dagger) - u(\pi_{x'} q^\dagger)\|_{L^2(\Omega)} \\ &\quad + \|z_n - u(q^\dagger)\|_{L^2(\Omega)} \leq Ch_n \|f\|_{\mathbb{H}^{1-s}(\Omega)} + \|u(q^\dagger) - u(\pi_{x'} q^\dagger)\|_{L^2(\Omega)} + \delta_n. \end{aligned} \quad (5.14)$$

We now invoke the extension property (2.15) and the identity (2.16) to arrive at

$$\begin{aligned} \|u(q^\dagger) - u(\pi_{x'} q^\dagger)\|_{L^2(\Omega)} &= \|\text{tr}_\Omega \mathcal{W}(q^\dagger) - \text{tr}_\Omega \mathcal{W}(\pi_{x'} q^\dagger)\|_{L^2(\Omega)} \\ &\leq \|\text{tr}_\Omega \mathcal{W}(q^\dagger) - \text{tr}_\Omega \mathcal{W}(\pi_{x'} q^\dagger)\|_{\mathbb{H}^s(\Omega)} = d_s^{-\frac{1}{2}} \|\mathcal{W}(q^\dagger) - \mathcal{W}(\pi_{x'} q^\dagger)\|_{\mathcal{C}}. \end{aligned} \quad (5.15)$$

It thus suffices to bound $\|\mathcal{W}(q^\dagger) - \mathcal{W}(\pi_{x'}q^\dagger)\|_{\mathcal{C}}$. Let $\phi \in \mathring{H}^1(y^\alpha, \mathcal{C})$. We invoke the problems that $\mathcal{W}(q^\dagger)$ and $\mathcal{W}(\pi_{x'}q^\dagger)$ solve to arrive at

$$\begin{aligned} a_{\mathcal{C}}(\mathcal{W}(q^\dagger) - \mathcal{W}(\pi_{x'}q^\dagger), \phi)(q^\dagger) &= \int_{\mathcal{C}} y^\alpha (\pi_{x'}q^\dagger - q^\dagger) \mathcal{W}(\pi_{x'}q^\dagger) \phi \, dx' \, dy \\ &\leq \|\pi_{x'}q^\dagger - q^\dagger\|_{L^\infty(\Omega)} \|\mathcal{W}(\pi_{x'}q^\dagger)\|_{L^2(y^\alpha, \mathcal{C})} \|\phi\|_{L^2(y^\alpha, \mathcal{C})}. \end{aligned}$$

This implies, in view of the estimate (5.10), that

$$\|\mathcal{W}(q^\dagger) - \mathcal{W}(\pi_{x'}q^\dagger)\|_{\mathcal{C}} \lesssim h_n^\gamma |q^\dagger|_{W^{\gamma, \infty}(\Omega)} \|f\|_{\mathbb{H}^{-s}(\Omega)}.$$

This estimate, combined with (5.14)–(5.15) allow us to conclude that

$$\|\operatorname{tr}_\Omega F(\pi_{x'}q^\dagger) - z_n\|_{L^2(\Omega)} \lesssim \delta_n + h_n^\gamma (|q^\dagger|_{W^{\gamma, \infty}(\Omega)} + 1) \|f\|_{\mathbb{H}^{1-s}(\Omega)}. \quad (5.16)$$

With the previous estimate at hand, we invoke (5.11)–(5.13) and conclude that

$$\limsup_{n \rightarrow \infty} \|Q_n - q^*\|_{L^2(\Omega)}^2 \leq \|q^\dagger - q^*\|_{L^2(\Omega)}^2. \quad (5.17)$$

The estimate (5.17) yields the existence of $\widehat{q} \in \mathcal{Q}$ and a non-relabeled subsequence $(Q_n)_{n \in \mathbb{N}}$ such that $(Q_n)_{n \in \mathbb{N}}$ converges weakly* to \widehat{q} in $L^\infty(\Omega)$. The arguments developed in the proof of Theorem 3.6 thus yield $\|\operatorname{tr}_\Omega(\mathcal{W}(Q_n) - \mathcal{W}(\widehat{q}))\|_{L^2(\Omega)} \rightarrow 0$ as $n \uparrow \infty$. On the other hand, the trace estimate (2.11) and (5.4), reveal that

$$\|\operatorname{tr}_\Omega(\mathcal{W}(Q_n) - F(Q_n))\|_{L^2(\Omega)} \lesssim \|\nabla(\mathcal{W}(Q_n) - F(Q_n))\|_{L^2(y^\alpha, \mathcal{C})} \lesssim h_n \|f\|_{\mathbb{H}^{1-s}(\Omega)} \rightarrow 0$$

as $n \uparrow \infty$. Consequently,

$$\begin{aligned} \|\operatorname{tr}_\Omega \mathcal{W}(\widehat{q}) - u^\dagger\|_{L^2(\Omega)} &= \lim_{n \rightarrow \infty} \|\operatorname{tr}_\Omega \mathcal{W}(Q_n) - z_n\|_{L^2(\Omega)} \\ &\leq \lim_{n \rightarrow \infty} \|\operatorname{tr}_\Omega(\mathcal{W}(Q_n) - F(Q_n))\|_{L^2(\Omega)} + \lim_{n \rightarrow \infty} \|\operatorname{tr}_\Omega F(Q_n) - z_n\|_{L^2(\Omega)} = 0, \end{aligned}$$

where we have used that

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{2} \|\operatorname{tr}_\Omega F(Q_n) - z_n\|_{L^2(\Omega)} &\leq \lim_{n \rightarrow \infty} \left(\frac{1}{2} \|\operatorname{tr}_\Omega F(\pi_{x'}q^\dagger) - z_n\|_{L^2(\Omega)} \right. \\ &\quad \left. + \frac{\rho_n}{2} \|\pi_{x'}q^\dagger - q^*\|_{L^2(\Omega)}^2 \right) = 0, \quad (5.18) \end{aligned}$$

which follows from (5.13), (5.14), and (5.16). We have thus concluded that $\widehat{q} \in \mathcal{I}(u^\dagger)$.

Finally, since q^\dagger solves (5.8), the fact that $(Q_n)_{n \in \mathbb{N}}$ converges weakly to \widehat{q} in $L^2(\Omega)$, and the estimate (5.17) allow us to conclude that

$$\begin{aligned} \|q^\dagger - q^*\|_{L^2(\Omega)}^2 &\leq \|\widehat{q} - q^*\|_{L^2(\Omega)}^2 \leq \liminf_{n \rightarrow \infty} \|Q_n - q^*\|_{L^2(\Omega)}^2 \\ &\leq \limsup_{n \rightarrow \infty} \|Q_n - q^*\|_{L^2(\Omega)}^2 \leq \|q^\dagger - q^*\|_{L^2(\Omega)}^2. \end{aligned}$$

Since the minimizer of (5.8) is unique, we conclude that $q^\dagger = \widehat{q}$ and that the whole sequence $(Q_n)_{n \in \mathbb{N}}$ converges to q^\dagger in the $L^2(\Omega)$ -norm. This concludes the proof. \square

5.3. A priori error estimates. In this section we provide an a priori error analysis for the fully discrete identification problem (5.6) when approximating solutions to (1.6). To accomplish this task, we follow [31] and introduce, for $\epsilon > 0$, $h_{\mathcal{T}} > 0$ and a local solution $r_{\delta,\rho}$ of problem (4.4), the following minimization problem:

$$\min_{Q \in \mathbb{Q}(r_{\delta,\rho})} \mathcal{D}_{\delta,\rho}(Q), \quad (5.19)$$

where the functional $\mathcal{D}_{\delta,\rho}$ is defined in (5.5) and $\mathbb{Q}(r_{\delta,\rho}) := \{Q \in \mathbb{Q}_{ad}(\mathcal{T}) : \|Q - r_{\delta,\rho}\|_{L^2(\Omega)} \leq \epsilon\}$. The next result guarantees existence of solutions for problem (5.19).

LEMMA 5.5 (existence of solutions). *If $h_{\mathcal{T}}$ is sufficiently small, then problem (5.19) has a solution $R_{\delta,\rho}$.*

Proof. Define $Q := \pi_{x'} r_{\delta,\rho}$. A density argument reveals that, if $h_{\mathcal{T}}$ is sufficiently small, then $\|Q - r_{\delta,\rho}\|_{L^2(\Omega)} \leq \epsilon$. Consequently, $\mathbb{Q}(r_{\delta,\rho}) \neq \emptyset$. We thus invoke standard arguments to obtain the desired result. This concludes the proof. \square

In the following result, we state a coercivity property for the second order derivatives of the discrete reduced cost functional in a neighborhood of a local solution $r_{\delta,\rho}$ of problem (4.4).

LEMMA 5.6 (local coercivity of $\mathcal{D}_{\delta,\rho}''$). *If $r_{\delta,\rho}$ denotes a local solution for problem (4.4) and $h_{\mathcal{T}}$ is sufficiently small, then there exist $\epsilon > 0$ such that for every $\hat{q} \in \mathbb{Q}$ that belongs to neighborhood $\|q - r_{\delta,\rho}\|_{L^2(\Omega)} \leq \epsilon$, we have that*

$$\mathcal{D}_{\delta,\rho}''(\hat{q})(q, q) \geq \frac{\theta}{4} \|q\|_{L^2(\Omega)}^2, \quad (5.20)$$

for all $q \in L^\infty(\Omega)$.

Proof. See [31, Lemma 4.7]. \square

LEMMA 5.7 (uniqueness). *Let $\epsilon > 0$ be sufficiently small such that (5.20) holds for $\hat{q} \in \mathbb{Q}(r_{\delta,\rho})$ and $q \in L^\infty(\Omega)$. If $h_{\mathcal{T}}$ is sufficiently small, then problem (5.19) admits a unique solution $R_{\delta,\rho}$.*

Proof. Let us assume that problem (5.19) admits two solutions R_1 and R_2 in $\mathbb{Q}(r_{\delta,\rho})$ which are such that $R_1 \neq R_2$. The differentiability properties of $\mathcal{D}_{\delta,\rho}$ yield

$$\mathcal{D}_{\delta,\rho}(R_1) = \mathcal{D}_{\delta,\rho}(R_2) + \mathcal{D}'_{\delta,\rho}(R_2)(R_1 - R_2) + \frac{1}{2} \mathcal{D}''_{\delta,\rho}(\tilde{R})(R_1 - R_2, R_1 - R_2),$$

where $\tilde{R} = \zeta R_1 + (1 - \zeta) R_2$ and $\zeta \in [0, 1]$. Notice that $\|\tilde{R} - r_{\delta,\rho}\|_{L^2(\Omega)} \leq \epsilon$.

Since R_2 solves (5.19), then $\mathcal{D}'_{\delta,\rho}(R_2)(R_1 - R_2) \geq 0$. This and an application of the second order optimality condition (5.20) allow us to conclude that

$$\mathcal{D}_{\delta,\rho}(R_1) \geq \mathcal{D}_{\delta,\rho}(R_2) + \frac{\theta}{4} \|R_1 - R_2\|_{L^2(\Omega)}^2,$$

which immediately yields $R_1 \equiv R_2$. This, that is a contradiction with the fact that $R_1 \neq R_2$, concludes the proof. \square

In the following result we show that the solution $R_{\delta,\rho}$ to problem (5.19) is a local solution to problem (5.6). To accomplish this task, we follow arguments elaborated in [15, 31]

LEMMA 5.8 ($R_{\delta,\rho}$ solves problem (5.6)). *Let $\epsilon > 0$ be sufficiently small such that (5.20) holds for $\hat{q} \in \mathbb{Q}(r_{\delta,\rho})$ and $q \in L^\infty(\Omega)$. Let $R_{\delta,\rho}$ be a solution of (5.19) such that $R_{\delta,\rho} \rightarrow r_{\delta,\rho}$ as $h \rightarrow 0$ in $L^2(\Omega)$. If $h_{\mathcal{T}}$ is sufficiently small, then $R_{\delta,\rho}$ is a solution of problem (5.6).*

Proof. Since $R_{\delta,\rho}$ solves (5.19), we have that

$$\mathcal{D}_{\delta,\rho}(R_{\delta,\rho}) \leq \mathcal{D}_{\delta,\rho}(Q) \quad \forall Q \in \mathbb{Q}_{ad}(\mathcal{T}) : \|Q - r_{\delta,\rho}\|_{L^2(\Omega)} \leq \epsilon. \quad (5.21)$$

Let $Q \in \mathbb{Q}_{ad}(\mathcal{T})$ such that $\|Q - R_{\delta,\rho}\|_{L^2(\Omega)} \leq \epsilon/2$. Then, since $R_{\delta,\rho} \rightarrow r_{\delta,\rho}$ as $h_{\mathcal{T}} \rightarrow 0$ in $L^2(\Omega)$, we have, for $h_{\mathcal{T}}$ sufficiently small, that

$$\|Q - r_{\delta,\rho}\|_{L^2(\Omega)} \leq \|Q - R_{\delta,\rho}\|_{L^2(\Omega)} + \|R_{\delta,\rho} - r_{\delta,\rho}\|_{L^2(\Omega)} \leq \epsilon/2 + \epsilon/2 = \epsilon.$$

Consequently, if $Q \in \mathbb{Q}_{ad}(\mathcal{T})$ is such that $\|Q - R_{\delta,\rho}\|_{L^2(\Omega)} \leq \epsilon/2$, then $\|Q - r_{\delta,\rho}\|_{L^2(\Omega)} \leq \epsilon$. In view of (5.21), we can thus conclude that

$$\mathcal{D}_{\delta,\rho}(R_{\delta,\rho}) \leq \mathcal{D}_{\delta,\rho}(Q) \quad \forall Q \in \mathbb{Q}_{ad}(\mathcal{T}) : \|Q - R_{\delta,\rho}\|_{L^2(\Omega)} \leq \epsilon/2,$$

This proves that $R_{\delta,\rho}$ solves (5.6) and concludes the proof. \square

We now derive a priori error estimates for our fully discrete scheme.

THEOREM 5.9 (a priori error estimates). *Let $r_{\delta,\rho}$ be a local solution for problem (4.4). Let ϵ and $h_{\mathcal{T}}$ sufficiently small such that the result of Lemma 5.7 hold. If $q^* \in H^1(\Omega)$, we thus have the following error estimate*

$$\|r_{\delta,\rho} - R_{\delta,\rho}\|_{L^2(\Omega)} \lesssim h_{\mathcal{T}},$$

where the hidden constant is independent of $r_{\delta,\rho}$, $R_{\delta,\rho}$, and $h_{\mathcal{T}}$.

Proof. We proceed in several steps.

1 We begin this step by choosing $\epsilon > 0$ sufficiently small such that, for $\hat{q} \in \mathcal{Q}$ that lies in the neighborhood $\|\hat{q} - r_{\delta,\rho}\|_{L^2(\Omega)} \leq \epsilon$, we have that

$$\mathcal{R}_{\delta,\rho}''(\hat{q})(q, q) \geq \frac{\theta}{2} \|q\|_{L^2(\Omega)}^2 \quad \forall q \in L^\infty(\Omega), \quad (5.22)$$

and, for $\hat{Q} \in \mathbb{Q}_{ad}(\mathcal{T})$ in the neighborhood $\|\hat{Q} - r_{\delta,\rho}\|_{L^2(\Omega)} \leq \epsilon$, we have that

$$\mathcal{D}_{\delta,\rho}''(\hat{Q})(q, q) \geq \frac{\theta}{4} \|q\|_{L^2(\Omega)}^2 \quad \forall q \in L^\infty(\Omega). \quad (5.23)$$

Let us now introduce the following optimization problem:

$$\min_{Q \in \mathbb{Q}(r_{\delta,\rho})} \mathcal{R}_{\delta,\rho}(Q), \quad (5.24)$$

We recall that $\mathbb{Q}(r_{\delta,\rho}) := \{Q \in \mathbb{Q}_{ad}(\mathcal{T}) : \|Q - r_{\delta,\rho}\|_{L^2(\Omega)} \leq \epsilon\}$ and that $\mathcal{R}_{\delta,\rho}$ is defined in (4.3). Notice that, if $h_{\mathcal{T}}$ is sufficiently small, then (5.24) has a unique solution $Z_{\delta,\rho}$.

A basic application of the triangle inequality allows us to arrive at

$$\|r_{\delta,\rho} - R_{\delta,\rho}\|_{L^2(\Omega)} \leq \|r_{\delta,\rho} - Z_{\delta,\rho}\|_{L^2(\Omega)} + \|Z_{\delta,\rho} - R_{\delta,\rho}\|_{L^2(\Omega)}. \quad (5.25)$$

2 We proceed to estimate the term $\|r_{\delta,\rho} - Z_{\delta,\rho}\|_{L^2(\Omega)}$. To accomplish this task, we set, for $\zeta \in [0, 1]$, $\hat{q} = \zeta r_{\delta,\rho} + (1 - \zeta)Z_{\delta,\rho}$ and notice that

$$\|\hat{q} - r_{\delta,\rho}\|_{L^2(\Omega)} = (1 - \zeta)\|r_{\delta,\rho} - Z_{\delta,\rho}\|_{L^2(\Omega)} \leq (1 - \zeta)\epsilon \leq \epsilon.$$

We can thus invoke (5.22) with $q = r_{\delta,\rho} - Z_{\delta,\rho}$ and obtain that

$$\begin{aligned} \frac{\theta}{2} \|r_{\delta,\rho} - Z_{\delta,\rho}\|_{L^2(\Omega)}^2 &\leq \mathcal{R}_{\delta,\rho}''(\hat{q})(r_{\delta,\rho} - Z_{\delta,\rho}, r_{\delta,\rho} - Z_{\delta,\rho}) \\ &= \mathcal{R}'_{\delta,\rho}(r_{\delta,\rho})(r_{\delta,\rho} - Z_{\delta,\rho}) - \mathcal{R}'_{\delta,\rho}(Z_{\delta,\rho})(r_{\delta,\rho} - Z_{\delta,\rho}) \\ &= \mathcal{R}'_{\delta,\rho}(r_{\delta,\rho})(r_{\delta,\rho} - Z_{\delta,\rho}) - \mathcal{R}'_{\delta,\rho}(Z_{\delta,\rho})(r_{\delta,\rho} - \pi_{x'} r_{\delta,\rho}) - \mathcal{R}'_{\delta,\rho}(Z_{\delta,\rho})(\pi_{x'} r_{\delta,\rho} - Z_{\delta,\rho}); \end{aligned}$$

$\pi_{x'}$ denotes the $L^2(\Omega)$ -orthogonal projection operator introduced in (5.9).

We now invoke the optimality condition (4.7) to arrive at $\mathcal{R}'_{\delta,\rho}(r_{\delta,\rho})(r_{\delta,\rho} - Z_{\delta,\rho}) \leq 0$. On the other hand, the optimality condition for problem (5.24), for $h_{\mathcal{T}}$ sufficiently small, yields $-\mathcal{R}'_{\delta,\rho}(Z_{\delta,\rho})(\pi_{x'}r_{\delta,\rho} - Z_{\delta,\rho}) \leq 0$. Thus,

$$\frac{\theta}{2} \|r_{\delta,\rho} - Z_{\delta,\rho}\|_{L^2(\Omega)}^2 \leq -\mathcal{R}'_{\delta,\rho}(Z_{\delta,\rho})(r_{\delta,\rho} - \pi_{x'}r_{\delta,\rho}).$$

Notice that $\mathcal{R}'_{\delta,\rho}(Z_{\delta,\rho})(r_{\delta,\rho} - \pi_{x'}r_{\delta,\rho}) = (\mathbf{r}(Z_{\delta,\rho}) + \rho(Z_{\delta,\rho} - q^*), r_{\delta,\rho} - \pi_{x'}r_{\delta,\rho})_{L^2(\Omega)}$. On the other hand, since $Z_{\delta,\rho} \in \mathbb{Q}(\mathcal{T})$, (5.9) yields $\rho(Z_{\delta,\rho}, r_{\delta,\rho} - \pi_{x'}r_{\delta,\rho})_{L^2(\Omega)} = 0$. We can thus use (5.9), again, to arrive at the estimate

$$\begin{aligned} \frac{\theta}{2} \|r_{\delta,\rho} - Z_{\delta,\rho}\|_{L^2(\Omega)}^2 &\leq \rho(q^* - \pi_{x'}q^*, r_{\delta,\rho} - \pi_{x'}r_{\delta,\rho})_{L^2(\Omega)} \\ &\quad - \rho(\mathbf{r}(Z_{\delta,\rho}) - \pi_{x'}\mathbf{r}(Z_{\delta,\rho}), r_{\delta,\rho} - \pi_{x'}r_{\delta,\rho})_{L^2(\Omega)}. \end{aligned}$$

Consequently,

$$\begin{aligned} \|r_{\delta,\rho} - Z_{\delta,\rho}\|_{L^2(\Omega)}^2 &\lesssim \|q^* - \pi_{x'}q^*\|_{L^2(\Omega)}^2 + \|r_{\delta,\rho} - \pi_{x'}r_{\delta,\rho}\|_{L^2(\Omega)}^2 \\ &+ \|\mathbf{r}(Z_{\delta,\rho}) - \pi_{x'}\mathbf{r}(Z_{\delta,\rho})\|_{L^2(\Omega)}^2 \lesssim h_{\mathcal{T}}^2 \left(\|\nabla q^*\|_{L^2(\Omega)}^2 + \|\nabla r_{\delta,\rho}\|_{L^2(\Omega)}^2 + \|\nabla \mathbf{r}(Z_{\delta,\rho})\|_{L^2(\Omega)}^2 \right). \end{aligned}$$

Notice that, in view of the fact that $q^* \in H^1(\Omega)$, the regularity results of Theorem 4.3 allow us to conclude that the norms involved in the right-hand side of the previous expression are uniformly bounded.

3 We now estimate $\|Z_{\delta,\rho} - R_{\delta,\rho}\|_{L^2(\Omega)}$ in (5.25). To accomplish this task, we set

$$\hat{Q} = \zeta R_{\delta,\rho} + (1 - \zeta)Z_{\delta,\rho}, \quad \zeta \in [0, 1],$$

and notice that $\|\hat{Q} - r_{\delta,\rho}\|_{L^2(\Omega)} \leq \zeta \|R_{\delta,\rho} - r_{\delta,\rho}\|_{L^2(\Omega)} + (1 - \zeta) \|Z_{\delta,\rho} - r_{\delta,\rho}\|_{L^2(\Omega)} \leq \epsilon$. We can thus invoke the second order optimality condition (5.23) with $q = Z_{\delta,\rho} - R_{\delta,\rho}$ to arrive at

$$\begin{aligned} \frac{\theta}{4} \|R_{\delta,\rho} - Z_{\delta,\rho}\|_{L^2(\Omega)}^2 &\leq \mathcal{D}''_{\delta,\rho}(\hat{Q})(R_{\delta,\rho} - Z_{\delta,\rho}, R_{\delta,\rho} - Z_{\delta,\rho}) \\ &= \mathcal{D}'_{\delta,\rho}(R_{\delta,\rho})(R_{\delta,\rho} - Z_{\delta,\rho}) - \mathcal{D}'_{\delta,\rho}(Z_{\delta,\rho})(R_{\delta,\rho} - Z_{\delta,\rho}). \end{aligned}$$

We now invoke the first-order optimality condition for problem (5.6) and conclude that $\mathcal{D}'_{\delta,\rho}(R_{\delta,\rho})(R_{\delta,\rho} - Z_{\delta,\rho}) \leq 0$. On the other hand, the first-order optimality condition for (5.24) yields $\mathcal{R}'_{\delta,\rho}(Z_{\delta,\rho})(R_{\delta,\rho} - Z_{\delta,\rho}) \geq 0$. Thus,

$$\frac{\theta}{4} \|R_{\delta,\rho} - Z_{\delta,\rho}\|_{L^2(\Omega)}^2 \leq \mathcal{R}'_{\delta,\rho}(Z_{\delta,\rho})(R_{\delta,\rho} - Z_{\delta,\rho}) - \mathcal{D}'_{\delta,\rho}(Z_{\delta,\rho})(R_{\delta,\rho} - Z_{\delta,\rho}).$$

Consequently, we can obtain that $\|R_{\delta,\rho} - Z_{\delta,\rho}\|_{L^2(\Omega)} \lesssim h_{\mathcal{T}}$.

4 The assertion follows from collecting all the estimates we obtained in previous steps. \square

We conclude with the following result.

COROLLARY 5.10 (a priori error estimates). *Let $r_{\delta,\rho}$ be a local solution for problem (4.4). Let ϵ and $h_{\mathcal{T}}$ sufficiently small such that the result of Lemma 5.7 hold. If*

$q^* \in H^1(\Omega)$, we thus have the existence of a sequence $\{Q_{\delta,\rho}\}$ of local minima of (5.6) such that

$$\|r_{\delta,\rho} - Q_{\delta,\rho}\|_{L^2(\Omega)} \lesssim h_{\mathcal{T}},$$

where the hidden constant is independent of $r_{\delta,\rho}$, $Q_{\delta,\rho}$, and $h_{\mathcal{T}}$.

Proof. The results of Lemma 5.8 show that $R_{\delta,\rho}$ solves (5.6). The desired error estimate thus follows from Theorem 5.9. \square

REFERENCES

- [1] M. Abramowitz and I.A. Stegun. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, volume 55 of *National Bureau of Standards Applied Mathematics Series*. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C., 1964.
- [2] L. Banjai, J.M. Melenk, R.H. Nochetto, E. Otárola, A.J. Salgado, and Ch. Schwab. Tensor FEM for spectral fractional diffusion. *arXiv:1707.07367v1*, 2017.
- [3] H.T. Banks and K. Kunisch. *Estimation Techniques for Distributed Parameter Systems*. Birkhauser Boston, 1989.
- [4] M. Benning and M. Burger. Modern regularization methods for inverse problems. *Acta Numer.*, 27:1–111, 2018.
- [5] M.Š. Birman and M.Z. Solomjak. *Spektralnaya teoriya samosopryazhennykh operatorov v gilbertovom prostranstve*. Leningrad. Univ., Leningrad, 1980.
- [6] C. Brändle, E. Colorado, A. de Pablo, and U. Sánchez. A concave-convex elliptic problem involving the fractional Laplacian. *Proc. Roy. Soc. Edinburgh Sect. A*, 143(1):39–71, 2013.
- [7] H. Brezis. *Functional analysis, Sobolev spaces and partial differential equations*. Universitext. Springer, New York, 2011.
- [8] C. Bucur and E. Valdinoci. *Nonlocal diffusion and applications*, volume 20 of *Lecture Notes of the Unione Matematica Italiana*. Springer, [Cham]; Unione Matematica Italiana, Bologna, 2016.
- [9] X. Cabré and Y. Sire. Nonlinear equations for fractional Laplacians II: Existence, uniqueness, and qualitative properties of solutions. *Trans. Amer. Math. Soc.*, 367(2):911–941, 2015.
- [10] X. Cabré and J. Tan. Positive solutions of nonlinear problems involving the square root of the Laplacian. *Adv. Math.*, 224(5):2052–2093, 2010.
- [11] L. Caffarelli and L. Silvestre. An extension problem related to the fractional Laplacian. *Comm. Part. Diff. Eqs.*, 32(7-9):1245–1260, 2007.
- [12] L. A. Caffarelli and A. Vasseur. Drift diffusion equations with fractional diffusion and the quasi-geostrophic equation. *Ann. of Math. (2)*, 171(3):1903–1930, 2010.
- [13] L.A. Caffarelli and P.R. Stinga. Fractional elliptic equations, Caccioppoli estimates and regularity. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 33(3):767–807, 2016.
- [14] A. Capella, J. Dávila, L. Dupaigne, and Y. Sire. Regularity of radial extremal solutions for some non-local semilinear equations. *Comm. Partial Differential Equations*, 36(8):1353–1384, 2011.
- [15] E. Casas and F. Tröltzsch. Error estimates for the finite-element approximation of a semilinear elliptic control problem. *Control Cybernet.*, 31(3):695–712, 2002. Well-posedness in optimization and related topics (Warsaw, 2001).
- [16] G. Chavent. *Nonlinear least squares for inverse problems*. Scientific Computation. Springer, New York, 2009. Theoretical foundations and step-by-step guide for applications.
- [17] F. Clarke. *Functional analysis, calculus of variations and optimal control*, volume 264 of *Graduate Texts in Mathematics*. Springer, London, 2013.
- [18] D. Colton and R. Kress. *Inverse Acoustic and Electromagnetic Scattering Theory*. Vol. 93 of Applied Mathematical Sciences, Springer, 2012.
- [19] J. Duoandikoetxea. *Fourier analysis*, volume 29 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2001. Translated and revised from the 1995 Spanish original by David Cruz-Uribe.
- [20] H. W. Engl, M. Hanke, and A. Neubauer. *Regularization of inverse problems*, volume 375 of *Mathematics and its Applications*. Kluwer Academic Publishers Group, Dordrecht, 1996.
- [21] A. Ern and J.-L. Guermond. *Theory and practice of finite elements*, volume 159 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 2004.
- [22] E.B. Fabes, C.E. Kenig, and R.P. Serapioni. The local regularity of solutions of degenerate elliptic equations. *Comm. Part. Diff. Eqs.*, 7(1):77–116, 1982.

- [23] G.B. Folland. *Real analysis*. Pure and Applied Mathematics (New York). John Wiley & Sons Inc., New York, second edition, 1999. Modern techniques and their applications, A Wiley-Interscience Publication.
- [24] D Fujiwara. Concrete characterization of the domains of fractional powers of some elliptic differential operators of the second order. *Proc. Japan Acad.*, 43:82–86, 1967.
- [25] V. Gol'dshtein and A. Ukhlov. Weighted Sobolev spaces and embedding theorems. *Trans. Amer. Math. Soc.*, 361(7):3829–3850, 2009.
- [26] P. Grisvard. *Elliptic problems in nonsmooth domains*, volume 69 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2011. Reprint of the 1985 original [MR0775683], With a foreword by Susanne C. Brenner.
- [27] D.N. Hao and T.N.T. Quyen. Convergence rates for tikhonov regularization of coefficient identification problems in laplace-type equations. *Inverse Problems*, 26(12):125014, 2010.
- [28] B. Kaltenbacher and A. Klassen. On convergence and convergence rates for ivanov and morozov regularization and application to some parameter identification problems in elliptic pdes. *Inverse Problems*, 34(5):055008, 2018.
- [29] T. Kato. *Perturbation theory for linear operators*. Classics in Mathematics. Springer-Verlag, Berlin, 1995. Reprint of the 1980 edition.
- [30] G. Kinderlehrer, D. Stampacchia. *An introduction to variational inequalities and their applications*, volume 88 of *Pure and Applied Mathematics*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1980.
- [31] A. Kröner and B. Vexler. A priori error estimates for elliptic optimal control problems with a bilinear state equation. *J. Comput. Appl. Math.*, 230(2):781–802, 2009.
- [32] A. Kufner and B. Opic. How to define reasonably weighted Sobolev spaces. *Comment. Math. Univ. Carolin.*, 25(3):537–554, 1984.
- [33] W. McLean. *Strongly elliptic systems and boundary integral equations*. Cambridge University Press, Cambridge, 2000.
- [34] G. Molica Bisci, V.D. Radulescu, and R. Servadei. *Variational methods for nonlocal fractional problems*, volume 162 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2016. With a foreword by Jean Mawhin.
- [35] B. Muckenhoupt. Weighted norm inequalities for the Hardy maximal function. *Trans. Amer. Math. Soc.*, 165:207–226, 1972.
- [36] R.H. Nochetto, E. Otárola, and A.J. Salgado. A PDE approach to fractional diffusion in general domains: a priori error analysis. *Found. Comput. Math.*, 15(3):733–791, 2015.
- [37] E. Otárola. *A PDE approach to numerical fractional diffusion*. PhD thesis, University of Maryland, College Park, 2014.
- [38] T.N.T. Quyen. Variational method for multiple parameter identification in elliptic pdes. *J. Math. Anal. Appl.*, 461(1):676–700, 2018.
- [39] T. Roubíček. *Nonlinear partial differential equations with applications*, volume 153 of *International Series of Numerical Mathematics*. Birkhäuser/Springer Basel AG, Basel, second edition, 2013.
- [40] T. Schuster, B. Kaltenbacher, B. Hofmann, and K.S. Kazimierski. *Regularization methods in Banach spaces*, volume 10 of *Radon Series on Computational and Applied Mathematics*. Walter de Gruyter GmbH & Co. KG, Berlin, 2012.
- [41] T. Schuster, B. Kaltenbacher, B. Hofmann, and K.S. Kazimierski. *Regularization Methods in Banach Spaces*, volume 10 of *Radon Series on Computational and Applied Mathematics*. de Gruyter, 2012.
- [42] P.R. Stinga and J.L. Torrea. Extension problem and Harnack's inequality for some fractional operators. *Comm. Partial Differential Equations*, 35(11):2092–2122, 2010.
- [43] A. Tarantola. *Inverse Problem Theory and Methods for Model Parameter Estimation*. SIAM, 2005.
- [44] F. Tröltzsch. *Optimal control of partial differential equations*, volume 112 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2010. Theory, methods and applications, Translated from the 2005 German original by Jürgen Sprekels.
- [45] B.O. Turesson. *Nonlinear potential theory and weighted Sobolev spaces*, volume 1736 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2000.
- [46] J.L. Vázquez. Recent progress in the theory of nonlinear diffusion with fractional Laplacian operators. *Discrete Contin. Dyn. Syst. Ser. S*, 7(4):857–885, 2014.