

1 **THE POISSON AND STOKES PROBLEMS ON WEIGHTED SPACES**  
2 **IN LIPSCHITZ DOMAINS AND UNDER SINGULAR FORCING\***

3 ENRIQUE OTÁROLA<sup>†</sup> AND ABNER J. SALGADO<sup>‡</sup>

4 **Abstract.** We show the well posedness of the Poisson and Stokes problems on weighted spaces  
5 over general Lipschitz domains. For a particular range of  $p$ , we consider those weights in the Muck-  
6 enhaupt class  $A_p$  that have no singularities in a neighborhood of the boundary of the domain.

7 **Key words.** Lipschitz domains, Muckenhoupt weights, weighted a priori estimates, elliptic  
8 equations, Stokes equations.

9 **AMS subject classifications.** 35D30, 35B45, 35J25.

10 **1. Introduction.** Let  $d \in \{2, 3\}$  and  $\Omega$  be a bounded domain of  $\mathbb{R}^d$  with Lip-  
11 schitz boundary  $\partial\Omega$ . Notice that we do not assume that  $\Omega$  is convex. The purpose  
12 of this work is to study the well posedness of the Dirichlet problem for the Poisson  
13 equation

14 (1) 
$$-\Delta u = F \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega,$$

15 and the Stokes problem

16 (2) 
$$-\Delta \mathbf{u} + \nabla \pi = -\operatorname{div} \mathbf{F}, \operatorname{div} \mathbf{u} = g, \text{ in } \Omega, \quad \mathbf{u} = 0 \text{ on } \partial\Omega,$$

17 where we allow the data  $F$  and  $(\mathbf{F}, g)$ , respectively to be singular.

18 The main technical tool that will allow us to assert certain degree of either reg-  
19 ularity or integrability on the singular data and solutions, is the theory of weighted  
20 spaces [20, 7]. This has been carried out with a large degree of success for smooth  
21 domains. On the other hand, to the best of our knowledge, in the case of, possibly  
22 convex, polytopes very little has been done in this direction. For instance, [6] proves a  
23 weighted Helmholtz decomposition on convex polytopes that is equivalent to the well  
24 posedness of (1). However, as described in [8], the argument presented there has a  
25 flaw. This was corrected in [8] for convex polytopes, and it is our intention here to, at  
26 least partially, remove the convexity assumption and study also the Stokes problem  
27 (2). We will obtain well posedness on weighted spaces, for a class of weights that do  
28 not have singularities or degeneracies near the boundary.

29 Our presentation will be organized as follows. Some preliminaries will be discussed  
30 in Section 2; where we will introduce the class of weights we shall operate with.  
31 The Poisson problem (1) will be studied in Section 3 along with some immediate  
32 applications of its well posedness. Finally, the Stokes problem (2) will be analyzed in  
33 Section 4.

34 **2. Preliminaries.** We will make repeated use of weighted Lebesgue and Sobolev  
35 spaces when the weight belongs to a Muckenhoupt class  $A_p$ . We refer the reader to  
36 [22, 21, 7, 13] for the basic facts about Muckenhoupt classes and the ensuing weighted

---

\*EO has been supported in part by CONICYT through FONDECYT project 3160201. AJS has been supported in part by NSF grant DMS-1720213.

<sup>†</sup>Departamento de Matemática, Universidad Técnica Federico Santa María, Valparaíso, Chile. ([enrique.otarola@usm.cl](mailto:enrique.otarola@usm.cl), <http://eotarola.mat.utfsm.cl/>).

<sup>‡</sup>Department of Mathematics, University of Tennessee, Knoxville, TN 37996, USA. ([asalgad1@utk.edu](mailto:asalgad1@utk.edu), <http://www.math.utk.edu/~abnersg>)

spaces. Here we only mention that a standard example of a Muckenhoupt weight is the distance to a lower dimensional object; see [2]. In particular, if  $z \in \Omega$  and we define the weight

$$(3) \quad \varpi_z(x) = |x - z|^\alpha,$$

then  $\varpi_z \in A_p$  provided that  $\alpha \in (-d, d(p-1))$ .

It is important to notice that in the example above, since  $z \in \Omega$ , there is a neighborhood of  $\partial\Omega$  where the weight  $\varpi_z$  has no degeneracies or singularities. In fact, it is continuous and strictly positive. This observation allows us to define a restricted class of Muckenhoupt weights for which our results will hold. The following definition is motivated by [9, Definition 2.5].

**DEFINITION 1** (class  $A_p(\Omega)$ ). *Let  $\Omega \subset \mathbb{R}^d$  be a Lipschitz domain. For  $p \in (1, \infty)$  we say that  $\varpi \in A_p$  belongs to  $A_p(\Omega)$  if there is an open set  $\mathcal{G} \subset \Omega$ , and positive constants  $\varepsilon > 0$  and  $\varpi_l > 0$  such that:*

1.  $\{x \in \Omega : \text{dist}(x, \partial\Omega) < \varepsilon\} \subset \mathcal{G}$ ,
2.  $\varpi \in C(\bar{\mathcal{G}})$ , and
3.  $\varpi_l \leq \varpi(x)$  for all  $x \in \bar{\mathcal{G}}$ .

We shall follow the convention that  $\omega$  will denote a weight in the class  $A_p$ , whereas  $\varpi$  one in the class  $A_p(\Omega)$ .

We shall also make use of the fact that if  $p \in (1, \infty)$ ,  $p' = p/(p-1)$  is its conjugate exponent, and  $\omega \in A_p$ , then  $\omega' := \omega^{-p'/p} \in A_{p'}$  with  $[\omega']_{A_{p'}} = [\omega]_{A_p}$ , where we set

$$[\omega]_{A_p} = \sup_B \left( \int_B \omega \right) \left( \int_B \omega' \right)^{p/p'}$$

and the supremum is taken over all balls  $B$ .

The ideas we will use to prove our well posedness results will, mainly, follow those used to prove [9, Theorem 5.2]. Essentially, owing to the fact that  $\varpi \in A_p(\Omega)$  is a regular function on a layer near the boundary of  $\Omega$ , we will use well posedness on weighted spaces for smooth domains in the interior and an unweighted result near the boundary and then patch these together. To be able to separate these two pieces we define cutoff functions  $\psi_i, \psi_\partial \in C_0^\infty(\mathbb{R}^d)$ ,  $\psi_i + \psi_\partial \equiv 1$  in  $\bar{\Omega}$  with the following properties:

- $\psi_i \equiv 1$  in a neighborhood of  $\Omega \setminus \mathcal{G}$ ,
- $\psi_i \equiv 0$  in a neighborhood of  $\partial\Omega$ , and
- setting  $\Omega_i$  to be the interior of  $\text{supp } \psi_i$ , then  $\partial\Omega_i \in C^{1,1}$ .

Note that, without loss of generality, we can assume that  $\partial\mathcal{G}$  is Lipschitz. Observe also that  $\text{supp } \nabla\psi_i \cup (\text{supp } \nabla\psi_\partial \cap \Omega) \subset \bar{\mathcal{G}}$ .

Finally, the relation  $A \lesssim B$  will mean that  $A \leq cB$  for a nonessential constant  $c$  that might change at each occurrence.

**3. The Poisson problem.** Let us now study problem (1). We begin by stating our definition of weak solution. Namely, for  $p \in (1, \infty)$  and  $\varpi \in A_p(\Omega)$ , given  $F \in W^{-1,p}(\varpi, \Omega)$  we seek for  $u \in W_0^{1,p}(\varpi, \Omega)$  such that

$$(4) \quad \int_\Omega \nabla u \nabla \varphi = \langle F, \varphi \rangle, \quad \forall \varphi \in C_0^\infty(\Omega).$$

Where by  $\langle \cdot, \cdot \rangle$  we denoted the duality pairing between  $W^{-1,p}(\varpi, \Omega)$  and  $W_0^{1,p'}(\varpi', \Omega)$ .

78 We will need two existence and uniqueness results for problem (4). The first one  
 79 deals with the well posedness of (4) on weighted spaces and  $C^1$  domains. For a proof  
 80 we refer the reader to [4, Theorem 2.5].

81 **THEOREM 2** (well posedness for  $C^1$  domains). *Let  $\Omega \subset \mathbb{R}^d$  be a bounded  $C^1$*   
 82 *domain,  $p \in (1, \infty)$  and  $\omega \in A_p$ . Then, for every  $F \in W^{-1,p}(\omega, \Omega)$  there is a unique*  
 83  *$u \in W_0^{1,p}(\omega, \Omega)$  that is a weak solution to (4) and, moreover, it satisfies*

$$84 \quad (5) \quad \|\nabla u\|_{\mathbf{L}^p(\omega, \Omega)} \lesssim \|F\|_{W^{-1,p}(\omega, \Omega)},$$

85 *where the hidden constant depends on  $\Omega$ ,  $[\omega]_{A_p}$ , and  $p$ , but it is independent of  $F$ .*

86 *Remark 3* (Theorem 2). Theorem 2 deserves the following comments:

- 87 • The definition of solution of (4) used in [4] assumes only that  $u \in W_0^{1,1}(\Omega)$ ; see the  
 88 statement of Theorem 2.5 in this reference. Under this assumption, the estimate  
 89 (5) of Theorem 2 (which is (2-13) of [4]) implies, using Conclusion i) of Corollary 1  
 90 of [10], that  $u \in W_0^{1,p}(\varpi, \Omega)$  so that our solutions coincide.
- 91 • [4, Theorem 2.5] assumes that (1) has a source term of the form  $F = -\operatorname{div} \mathbf{f}$  with  
 92  $\mathbf{f} \in \mathbf{L}^p(\varpi, \Omega)$ . However, as we will do below in Corollary 9, from such a result  
 93 inf-sup conditions, and consequently well posedness, can be derived.

94 The second result deals with the well posedness of (4) on Lipschitz domains. This  
 95 result can be found in [15, Theorem 2] and [16, Theorem 0.5].

96 **THEOREM 4** (well posedness for Lipschitz domains). *Let  $\Omega \subset \mathbb{R}^d$  be a bounded*  
 97 *Lipschitz domain. There exists*

$$98 \quad (6) \quad p_1 > \begin{cases} 3 & d = 3, \\ 4 & d = 2, \end{cases}$$

99 *depending solely on the Lipschitz constant of  $\partial\Omega$  such that, if  $p_0 = p'_1$ , and  $p \in (p_0, p_1)$ ,*  
 100 *then for every  $F \in W^{-1,p}(\Omega)$  there is a unique  $u \in W_0^{1,p}(\Omega)$  that is a weak solution*  
 101 *to (4) and, moreover, it satisfies*

$$102 \quad \|\nabla u\|_{\mathbf{L}^p(\Omega)} \lesssim \|F\|_{W^{-1,p}(\Omega)},$$

103 *where the hidden constant depends on  $\Omega$ , and  $p$ , but it is independent of  $F$ .*

104 We are now in position to state the well posedness of (4).

105 **THEOREM 5** (well posedness on weighted spaces for Lipschitz domains). *Let*  
 106  *$\Omega \subset \mathbb{R}^d$  be a bounded Lipschitz domain. There is  $p_1$  satisfying (6), such that, if  $p_0 = p'_1$ ,*  
 107  *$p \in (p_0, p_1)$ , and  $\varpi \in A_p(\Omega)$ . Then, for every  $F \in W^{-1,p}(\varpi, \Omega)$  there is a unique*  
 108  *$u \in W_0^{1,p}(\varpi, \Omega)$  that is a weak solution to (4) and, moreover, it satisfies*

$$109 \quad (7) \quad \|\nabla u\|_{\mathbf{L}^p(\varpi, \Omega)} \lesssim \|F\|_{W^{-1,p}(\varpi, \Omega)},$$

110 *where the hidden constant depends on  $\Omega$ ,  $[\varpi]_{A_p}$ , and  $p$ , but it is independent of  $F$ .*

111 Before proving this result, we first establish a preliminary a priori estimate.

112 **LEMMA 6** (Gårding-like inequality). *Let  $\Omega$ ,  $p$  and  $\varpi$  be as in Theorem 5. If*  
 113  *$u \in W_0^{1,p}(\varpi, \Omega)$  is a weak solution of (4), then it satisfies*

$$114 \quad \|\nabla u\|_{\mathbf{L}^p(\varpi, \Omega)} \lesssim \|F\|_{W^{-1,p}(\varpi, \Omega)} + \|u\|_{L^p(\mathcal{G})},$$

115 *where the hidden constant depends on  $\mathcal{G}$ ,  $p$  and  $[\varpi]_{A_p}$ , but it is independent of  $F$ .*

116 *Proof.* Let  $u_i = u\psi_i \in W_0^{1,p}(\varpi, \Omega_i)$  and  $\varphi \in C_0^\infty(\Omega_i)$  then

$$117 \quad (8) \quad \begin{aligned} \int_{\Omega_i} \nabla u_i \nabla \varphi &= \int_{\Omega_i} \nabla u \nabla (\psi_i \varphi) - \int_{\Omega_i} \varphi \nabla u \nabla \psi_i + \int_{\Omega_i} u \nabla \psi_i \nabla \varphi \\ &= \int_{\Omega_i} \nabla u \nabla (\psi_i \varphi) + \int_{\mathcal{G}} u \operatorname{div}(\varphi \nabla \psi_i) + \int_{\mathcal{G}} u \nabla \psi_i \nabla \varphi, \end{aligned}$$

118 where we used that  $\operatorname{supp} \nabla \psi_i \subset \bar{\mathcal{G}}$ . This identity shows that  $u_i$  is a weak solution to  
119 (4) over  $\Omega_i \in C^{1,1}$  with right hand side  $F_i$  defined by

$$120 \quad \langle F_i, \varphi \rangle := \langle F, \psi_i \varphi \rangle + \int_{\mathcal{G}} u \operatorname{div}(\varphi \nabla \psi_i) + \int_{\mathcal{G}} u \nabla \psi_i \nabla \varphi.$$

121 Consequently, invoking the estimate of Theorem 2 we can obtain that

$$122 \quad \|\nabla u_i\|_{\mathbf{L}^p(\varpi, \Omega_i)} \lesssim \|F_i\|_{W^{-1,p}(\varpi, \Omega_i)}.$$

123 Now, using the fact that  $\varpi$ , when restricted to  $\mathcal{G}$  is uniformly positive and bounded  
124 we can estimate

$$125 \quad \begin{aligned} \|F_i\|_{W^{-1,p}(\varpi, \Omega_i)} &\lesssim \|F\|_{W^{-1,p}(\varpi, \Omega)} + \sup_{0 \neq \varphi \in W_0^{1,p'}(\varpi', \Omega_i)} \frac{\int_{\mathcal{G}} |u| |\nabla \varphi|}{\|\nabla \varphi\|_{\mathbf{L}^{p'}(\varpi', \Omega_i)}} \\ 126 \quad &+ \sup_{0 \neq \varphi \in W_0^{1,p'}(\varpi', \Omega_i)} \frac{\int_{\mathcal{G}} |u| |\varphi|}{\|\nabla \varphi\|_{\mathbf{L}^{p'}(\varpi', \Omega_i)}} \\ 127 \quad &\lesssim \|F\|_{W^{-1,p}(\varpi, \Omega)} + \|u\|_{L^p(\mathcal{G})}. \end{aligned}$$

129 Combining the previous two bounds allows us to conclude

$$130 \quad (9) \quad \|\nabla u_i\|_{\mathbf{L}^p(\varpi, \Omega_i)} \lesssim \|F\|_{W^{-1,p}(\varpi, \Omega)} + \|u\|_{L^p(\mathcal{G})}.$$

131 Define now  $u_\partial = u\psi_\partial \in W_0^{1,p}(\mathcal{G})$ . Similar computations, but using now Theorem 4  
132 for the Lipschitz domain  $\mathcal{G}$  allow us to conclude

$$133 \quad \|\nabla u_\partial\|_{\mathbf{L}^p(\mathcal{G})} \lesssim \|F\|_{W^{-1,p}(\varpi, \Omega)} + \|u\|_{L^p(\mathcal{G})}$$

134 so that, using the uniform boundedness and positivity of  $\varpi$  over  $\mathcal{G}$  we conclude

$$135 \quad (10) \quad \|\nabla u_\partial\|_{\mathbf{L}^p(\varpi, \mathcal{G})} \lesssim \|F\|_{W^{-1,p}(\varpi, \Omega)} + \|u\|_{L^p(\mathcal{G})}.$$

136 Since  $u = u_i + u_\partial$ , an application of the triangle inequality, and estimates (9) and  
137 (10) yield the desired bound.  $\square$

138 We are now in position to begin proving Theorem 5 with the uniqueness result.

139 **LEMMA 7 (uniqueness).** *Let  $\Omega \subset \mathbb{R}^d$  be a bounded Lipschitz domain. There is*  
140  *$p_1$  satisfying (6) such that, whenever  $p \in [2, p_1)$ , and  $\varpi \in A_p(\Omega)$  we have that if*  
141  *$u \in W_0^{1,p}(\varpi, \Omega)$  solves (4) with  $F = 0$ , then  $u = 0$ .*

142 *Proof.* We begin by observing that the assumptions imply that  $u$  is a solution of  
143  $-\Delta u = 0$  in  $\mathcal{D}'(\Omega_i)$ . Thus, we obtain that  $u \in W^{2,r}(\Omega_i)$  for every  $r \in (1, \infty)$ , [12,  
144 Theorem 9.15]; notice that  $\partial\Omega_i \in C^{1,1}$ . Further, similar computations to the ones  
145 that led to (8) reveal that, for all  $\varphi \in C_0^\infty(\Omega_i)$ , we have

$$146 \quad \left| \int_{\Omega_i} \nabla u_i \nabla \varphi \right| \lesssim \|\nabla \varphi\|_{\mathbf{L}^{r'}(\Omega_i)}$$

147 where the hidden constant depends on  $r$  and  $u$ . This shows that  $\varphi \mapsto \int_{\Omega_i} \nabla u_i \nabla \varphi$   
 148 defines an element of  $W^{-1,r}(\Omega_i)$  so that, by Theorem 4, we obtain that  $u_i \in W_0^{1,2}(\Omega_i)$ .

149 Since we are assuming that  $\varpi \in A_p(\Omega)$ , and,  $p \geq 2$ , we also have that  $u_\partial \in$   
 150  $W_0^{1,p}(\varpi, \mathcal{G}) = W_0^{1,p}(\mathcal{G}) \hookrightarrow W_0^{1,2}(\mathcal{G})$  so that, to conclude

$$151 \quad u = u_i + u_\partial \in W_0^{1,2}(\Omega).$$

152 This allows us to set  $\varphi = u$  in the condition to obtain that  $\nabla u = 0$  almost everywhere  
 153 and, thus,  $u = 0$ .  $\square$

154 *Remark 8* (alternative proof). Uniqueness can also be obtained as follows. Since  
 155  $u \in W_0^{1,p}(\varpi, \Omega) \subset W_0^{1,1}(\Omega)$  then we have, in particular, that  $u \in L^1(\Omega)$  and that

$$156 \quad \int_{\Omega} u \Delta \varphi = 0 \quad \forall \varphi \in C_0^\infty(\Omega).$$

157 Now, from this we infer that  $u$  is a.e. equal to a  $C^2(\Omega)$  and harmonic function. To  
 158 see this, we note that, if  $\rho_\epsilon$  is a radial mollifier, then for  $\epsilon$  sufficiently small we have  
 159 that  $\varphi \star \rho_\epsilon \in C_0^\infty(\Omega)$  and, thus,

$$160 \quad \int (u \star \rho_\epsilon) \Delta \varphi = \int u \Delta (\varphi \star \rho_\epsilon) = 0.$$

161 Since  $u \star \rho_\epsilon \in C(\Omega)$ , we can then invoke [14, Theorem 1.16] to conclude that  $u \star \rho_\epsilon$  is  
 162 harmonic in  $\Omega$ . This, by [14, Theorem 1.6] implies that  $u \star \rho_\epsilon$  satisfies the mean value  
 163 property

$$164 \quad u \star \rho_\epsilon(x) = \int_{B_r(x)} u \star \rho_\epsilon = \int_{B_R(x)} u \star \rho_\epsilon \quad \forall x \in \Omega, \quad B_r(x), B_R(x) \subset \Omega.$$

165 Define, for all  $x \in \Omega$  and any  $r$  such that  $B_r(x) \subset \Omega$

$$\bar{u}(x) = \int_{B_r(x)} u.$$

166 Notice that  $\bar{u}$  is continuous,  $u \star \rho_\epsilon \rightarrow \bar{u}$  for every  $x \in \Omega$  and in  $L^1_{loc}(\Omega)$ , and  $u = \bar{u}$   
 167 almost everywhere. Since  $\bar{u}$  satisfies the mean value property, then [14, Theorem 1.8]  
 168 yields that  $\bar{u} \in C^2(\Omega)$  and is harmonic. As a consequence  $u_i = u\psi_i \in W_0^{1,2}(\Omega)$ .  $\blacksquare$

170 We thank the anonymous reviewer for suggesting this alternative proof.

171 Having shown uniqueness we can finally prove Theorem 5.

172 *Proof of Theorem 5.* Consider first  $p \in [2, p_1)$  and assume that (7) is false. If that  
 173 is the case, then it is possible to find sequences  $(u_k, F_k) \in W_0^{1,p}(\varpi, \Omega) \times W^{-1,p}(\varpi, \Omega)$   
 174 such that they satisfy (4) with  $\|\nabla u_k\|_{\mathbf{L}^p(\varpi, \Omega)} = 1$ , but  $F_k \rightarrow 0$  in  $W^{-1,p}(\varpi, \Omega)$ , as  
 175  $k \rightarrow \infty$ . By passing to a, not relabeled, subsequence we can assume that  $u_k \rightharpoonup u \in$   
 176  $W_0^{1,p}(\varpi, \Omega)$  and that this limit satisfies (4) for  $F = 0$ , so that, by Lemma 7, we have  
 177 that  $u = 0$ . On the other hand, the compact embedding of  $W_0^{1,p}(\varpi, \Omega)$  into  $L^p(\varpi, \Omega)$   
 178 shows that  $u_k \rightarrow 0$  in  $L^p(\varpi, \Omega)$ , so that  $\|u\|_{L^p(\mathcal{G})} = 0$ . Consequently, using Lemma 6,  
 179 we have that

$$180 \quad 1 = \|\nabla u_k\|_{\mathbf{L}^p(\varpi, \Omega)} \lesssim \|F_k\|_{W^{-1,p}(\varpi, \Omega)} + \|u_k\|_{L^p(\mathcal{G})} \rightarrow 0, \quad k \uparrow \infty,$$

181 which is a contradiction.

182 With the a priori estimate (7) at hand we can now show existence of a solution  
 183  $u \in W_0^{1,p}(\varpi, \Omega)$ , in the case  $p \in [2, p_1)$ , by an approximation argument. Indeed,  
 184 given  $F \in W^{-1,p}(\varpi, \Omega)$  we construct a sequence  $F_k \in C^\infty(\Omega)$  such that  $F_k \rightarrow F$  in  
 185  $W^{-1,p}(\varpi, \Omega)$ . Theorem 4 then guarantees the existence of a unique  $u_k \in W_0^{1,p}(\Omega)$   
 186 that solves (4) with right hand side  $F_k$ . To be able to pass to the limit with (7) it is  
 187 then necessary to show that  $u_k \in W_0^{1,p}(\varpi, \Omega)$ :

- 188 • Since  $\varpi \in A_p(\Omega)$ , then  $u_k \in W^{1,p}(\varpi, \mathcal{G})$ .
- 189 • Since  $\varpi \in A_p$ , we invoke the *reverse Hölder inequality* [7, Theorem 5.4], and  
 190 conclude the existence of  $\gamma > 0$  such that  $\varpi^{1+\gamma} \in L^1(\Omega_i)$ . Now, given that  
 191  $F_k \in C^\infty(\Omega)$ , we can invoke [12, Theorem 8.10] to obtain that  $u_k \in W^{r,2}(\Omega_i)$   
 192 with  $r$  so large that, by Sobolev embedding, the right hand side of the inequality

$$193 \quad \int_{\Omega_i} \varpi |\nabla u_k|^p \leq \left( \int_{\Omega_i} \varpi^{1+\gamma} \right)^{1/(1+\gamma)} \left( \int_{\Omega_i} |\nabla u_k|^{p(1+\gamma)/\gamma} \right)^{\gamma/(1+\gamma)}$$

194 is finite.

195 This shows that  $u_k \in W_0^{1,p}(\varpi, \Omega)$  and, thus, existence of a solution.

196 Having proved the result for  $p \in [2, p_1)$ , the assertion for  $p \in (p_0, 2)$  follows by  
 197 duality.  $\square$

198 **3.1. Application. Well posedness with Dirac sources.** Let us discuss some  
 199 applications of our main result. An immediate corollary is the following.

200 **COROLLARY 9** (inf-sup condition). *Let  $\Omega \subset \mathbb{R}^d$  be a bounded Lipschitz domain.*  
 201 *There is  $p_1$ , depending solely on the Lipschitz constant of  $\partial\Omega$ , that satisfies (6), and*  
 202 *such that, if  $p_0 = p'_1$ ,  $p \in (p_0, p_1)$ , and  $\varpi \in A_p(\Omega)$ , we thus have, for every  $v \in$   
 203  $W_0^{1,p}(\varpi, \Omega)$ , that*

$$204 \quad \|\nabla v\|_{\mathbf{L}^p(\varpi, \Omega)} \lesssim \sup_{0 \neq w \in W_0^{1,p'}(\varpi', \Omega)} \frac{\int_{\Omega} \nabla v \nabla w}{\|\nabla w\|_{\mathbf{L}^{p'}(\varpi', \Omega)}}$$

205 where the hidden constant is independent of  $v$ .

206 *Proof.* Given  $v \in W_0^{1,p}(\varpi, \Omega)$  we observe that  $\varpi |\nabla v|^{p-2} \nabla v \in \mathbf{L}^{p'}(\varpi', \Omega)$  so that  
 207 the functional  $F_v = -\operatorname{div}(\varpi |\nabla v|^{p-2} \nabla v) \in W^{-1,p'}(\varpi', \Omega)$  with

$$208 \quad \|F_v\|_{W^{-1,p'}(\varpi', \Omega)} \lesssim \|\nabla v\|_{\mathbf{L}^p(\varpi, \Omega)}^{p-1}.$$

209 By Theorem 5 there is a unique function  $w_v \in W_0^{1,p'}(\varpi', \Omega)$  that solves (4) with right  
 210 hand side  $F_v$ , i.e.,

$$211 \quad \int_{\Omega} \nabla w_v \nabla \varphi = \int_{\Omega} \varpi |\nabla v|^{p-2} \nabla v \nabla \varphi \quad \forall \varphi \in W_0^{1,p}(\varpi, \Omega),$$

212 with the corresponding estimate. Thus, setting  $\varphi = v$  the assertion follows.  $\square$

213 The inf-sup condition of Corollary 9 allows us to then establish the well posedness  
 214 of the Poisson problem with Dirac sources on weighted spaces.

215 **COROLLARY 10** (well posedness). *Let  $\Omega \subset \mathbb{R}^d$ , with  $d \in \{2, 3\}$ , be a bounded Lip-*  
 216 *schitz domain and  $z \in \Omega$ . Then, for  $\alpha \in (d-2, d)$ , and  $\varpi_z$  defined as in (3), there is*  
 217 *a unique  $u \in W_0^{1,2}(\varpi_z, \Omega)$  that is a weak solution of*

$$218 \quad -\Delta u = \delta_z \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega.$$

219 *Proof.* Notice that, since  $\alpha \in (d-2, d) \subset (-d, d)$  and  $z \in \Omega$ , we have that  $\varpi_z \in$   
 220  $A_2(\Omega)$ . In light of Corollary 9 we only need to prove then that  $\delta_z \in W^{-1,2}(\varpi_z, \Omega)$ ,  
 221 but this follows from [17, Lemma 7.1.3] when  $\alpha \in (d-2, d)$ ; see also [1, Theorem 2.3].  
 222 This concludes the proof.  $\square$

223 **3.2. A weighted Helmholtz decomposition on Lipschitz domains.** As  
 224 the results of [9, 10] show, in the study of the Stokes problem (2) it is sometimes  
 225 necessary to have a weighted decomposition of the spaces  $\mathbf{L}^p(\varpi, \Omega)$ , where the weight  
 226 is adapted to the singularity of  $\mathbf{F}$ . Here we show such a decomposition for a Lipschitz  
 227 domain and for a weight of class  $A_p(\Omega)$ .

228 We introduce some notation. For  $p \in (1, \infty)$  and a weight  $\varpi \in A_p(\Omega)$ , the space  
 229 of solenoidal functions is

$$230 \quad \mathbf{L}_{\sigma, N}^p(\varpi, \Omega) = \{\mathbf{v} \in \mathbf{L}^p(\varpi, \Omega) : \operatorname{div} \mathbf{v} = 0\}.$$

231 The space of gradients is

$$232 \quad \mathbf{G}_D^p(\varpi, \Omega) = \left\{ \nabla v : v \in W_0^{1,p}(\varpi, \Omega) \right\}.$$

233 We wish to show the decomposition

$$234 \quad (11) \quad \mathbf{L}^p(\varpi, \Omega) = \mathbf{L}_{\sigma, N}^p(\varpi, \Omega) \oplus \mathbf{G}_D^p(\varpi, \Omega)$$

235 with a continuous projection  $\mathcal{P}_{p, \varpi} : \mathbf{L}^p(\varpi, \Omega) \rightarrow \mathbf{L}_{\sigma, N}^p(\varpi, \Omega)$  such that  $\ker \mathcal{P}_{p, \varpi} =$   
 236  $\mathbf{G}_D^p(\varpi, \Omega)$ .

237 **COROLLARY 11** (weighted Helmholtz decomposition I). *Let  $\Omega$ ,  $p_1$ ,  $p$  and  $\varpi$  be*  
 238 *as in Theorem 5. Then, the decomposition (11) holds.*

239 *Proof.* Let  $\mathbf{f} \in \mathbf{L}^p(\varpi, \Omega)$ . By Theorem 5 there is a unique  $u \in W_0^{1,p}(\varpi, \Omega)$  that  
 240 solves (4) with  $F = \operatorname{div} \mathbf{f}$ . Setting  $\mathbf{f} = (\mathbf{f} - \nabla u) + \nabla u$  gives, by uniqueness and the  
 241 estimate on  $\nabla u$ , the desired decomposition.  $\square$

242 **3.3. Variable coefficients.** We conclude the discussion on the Dirichlet prob-  
 243 lem (1) by showing how, from Theorem 5, we can assert the well posedness of a  
 244 problem with variable coefficients, thus obtaining a weighted version of Meyers' result  
 245 [18]. Namely, let  $\mathcal{A} \in \mathbf{L}^\infty(\Omega)$  be a matrix-valued coefficient such that:

- 246 • For almost every  $x \in \Omega$ ,  $\mathcal{A}(x)$  is symmetric,
- 247 • There are constants  $\lambda, \Lambda \in \mathbb{R}$  with  $0 < \lambda \leq \Lambda$  such that, for almost every  $x \in \Omega$ ,

$$248 \quad \lambda |\boldsymbol{\xi}|^2 \leq \boldsymbol{\xi}^\top \mathcal{A}(x) \boldsymbol{\xi} \leq \Lambda |\boldsymbol{\xi}|^2 \quad \forall \boldsymbol{\xi} \in \mathbb{R}^d,$$

249 where  $|\cdot|$  denotes the Euclidean norm in  $\mathbb{R}^d$ .

250 Let  $\Omega \subset \mathbb{R}^d$  be a bounded Lipschitz domain,  $p \in (1, \infty)$ , and  $\varpi \in A_p(\Omega)$ . Given  
 251  $F \in W^{-1,p}(\varpi, \Omega)$ , the purpose of this section is to study the well posedness of the  
 252 following problem: find  $v \in W_0^{1,p}(\Omega)$  such that

$$253 \quad (12) \quad \int_{\Omega} \nabla \varphi^\top \mathcal{A} \nabla v = \langle F, \varphi \rangle \quad \forall \varphi \in C_0^\infty(\Omega).$$

254 As it is well known, even in the unweighted case, problem (12) is not generally  
 255 well posed for  $p \neq 2$ . This heavily depends on the behavior of  $\mathcal{A}$ ; see [18]. More  
 256 specifically it depends on the quantity

$$257 \quad (13) \quad \varrho(\mathcal{A}) = \frac{\lambda}{\Lambda}.$$

258 The following result is inspired by [3, Proposition 1].

259 THEOREM 12 (well posedness with variable coefficients for Lipschitz domains).  
 260 Let  $\Omega \subset \mathbb{R}^d$  be a bounded Lipschitz domain, and  $p$  and  $\varpi$  be as in Theorem 5. There  
 261 is  $\varrho_0$  such that, if  $\varrho(\mathcal{A}) > \varrho_0$ , the problem (12) is well posed and it has the estimate

$$262 \quad \|\nabla v\|_{\mathbf{L}^p(\varpi, \Omega)} \lesssim \|F\|_{W^{-1,p}(\varpi, \Omega)},$$

263 where the hidden constant depends on  $\Omega$ ,  $p$ ,  $[\varpi]_{A_p}$  and  $\varrho(\mathcal{A})$ , but it is independent of  
 264  $F$ .

265 *Proof.* For  $p$  in the indicated range, Theorem 5 shows that the mapping  $T :=$   
 266  $-\Delta : W_0^{1,p}(\varpi, \Omega) \rightarrow W^{-1,p}(\varpi, \Omega)$  is invertible. In other words, there is a constant  
 267  $C(\Delta, p, \varpi)$  such that

$$268 \quad \|T^{-1}\|_{\mathcal{L}(W^{-1,p}(\varpi, \Omega), W_0^{1,p}(\varpi, \Omega))} \leq C(\Delta, p, \varpi).$$

269 Define  $S : W_0^{1,p}(\varpi, \Omega) \rightarrow W^{-1,p}(\varpi, \Omega)$  via

$$270 \quad \langle Sw, \varphi \rangle = \int_{\Omega} \frac{1}{\Lambda} \nabla \varphi^\top \mathcal{A} \nabla w.$$

271 Notice that

$$272 \quad \|Sw\|_{W^{-1,p}(\varpi, \Omega)} \leq \frac{1}{\Lambda} \|\mathcal{A} \nabla w\|_{\mathbf{L}^p(\varpi, \Omega)} \leq \|\nabla w\|_{\mathbf{L}^p(\varpi, \Omega)},$$

273 which implies

$$274 \quad \|S\|_{\mathcal{L}(W_0^{1,p}(\varpi, \Omega), W^{-1,p}(\varpi, \Omega))} \leq 1.$$

275 Let now  $Q = T - S : W_0^{1,p}(\varpi, \Omega) \rightarrow W^{-1,p}(\varpi, \Omega)$  and notice that

$$276 \quad \langle Qw, \varphi \rangle = \int_{\Omega} \nabla \varphi^\top \left( \mathcal{I} - \frac{1}{\Lambda} \mathcal{A} \right) \nabla w,$$

277 where  $\mathcal{I}$  is the identity matrix. This implies that

$$278 \quad \|Q\|_{\mathcal{L}(W_0^{1,p}(\varpi, \Omega), W^{-1,p}(\varpi, \Omega))} = \left\| \max \left\{ \lambda : \lambda \in \sigma \left( \mathcal{I} - \frac{1}{\Lambda} \mathcal{A} \right) \right\} \right\|_{\mathbf{L}^\infty(\Omega)}.$$

279 But, the conditions on  $\mathcal{A}$  imply that, for almost every  $x \in \Omega$ ,

$$280 \quad \lambda \mathcal{I} \preceq \mathcal{A}(x) \preceq \Lambda \mathcal{I} \quad \implies \quad 0 \preceq \mathcal{I} - \frac{1}{\Lambda} \mathcal{A}(x) \preceq (1 - \varrho(\mathcal{A})) \mathcal{I},$$

281 where  $\preceq$  means an inequality in the spectral sense. From this we conclude that

$$282 \quad \max \left\{ \lambda : \lambda \in \sigma \left( \mathcal{I} - \frac{1}{\Lambda} \mathcal{A} \right) \right\} \leq 1 - \varrho(\mathcal{A}).$$

283 We have now that

$$284 \quad \|T^{-1}Q\|_{\mathcal{L}(W_0^{1,p}(\varpi, \Omega))} \leq C(\Delta, p, \varpi)(1 - \varrho(\mathcal{A})),$$

285 and, since  $S = T - Q = T(I - T^{-1}Q)$ , we have that  $S$  is invertible, provided  
 286  $C(\Delta, p, \varpi)(1 - \varrho(\mathcal{A})) < 1$  which holds if

$$287 \quad \varrho(\mathcal{A}) > \varrho_0 = 1 - \frac{1}{C(\Delta, p, \varpi)}.$$



288 If that is the case, then

$$289 \quad \|S^{-1}\|_{\mathcal{L}(W^{-1,p}(\varpi,\Omega), W_0^{1,p}(\varpi,\Omega))} \leq \frac{C(\Delta, p, \varpi)}{1 - C(\Delta, p, \varpi)(1 - \varrho(\mathcal{A}))},$$

290 which by linearity implies that (12) has a unique solution with the estimate

$$291 \quad \|\nabla v\|_{\mathbf{L}^p(\varpi,\Omega)} \leq \frac{1}{\Lambda} \frac{C(\Delta, p, \varpi)}{1 - C(\Delta, p, \varpi)(1 - \varrho(\mathcal{A}))} \|F\|_{W^{-1,p}(\varpi,\Omega)}.$$

292 The theorem is thus proved.  $\square$

293 **3.4. The Neumann problem.** We briefly comment that, with the same tech-  
 294 niques, our result can be transferred to the case of Neumann boundary conditions.  
 295 For that, all that is needed is the analogues to Theorems 2 and 4 to carry out our  
 296 considerations.

297 **THEOREM 13** (well posedness of the Neumann problem in Lipschitz domains).  
 298 *Let  $\Omega \subset \mathbb{R}^d$  be a bounded Lipschitz domain. There is  $p_1$  that satisfies (6), such that if*  
 299  *$p_0 = p'_1$ ,  $p \in (p_0, p_1)$ , and  $\varpi \in A_p(\Omega)$ . then, for every  $\mathbf{f} \in \mathbf{L}^p(\varpi, \Omega)$  there is a unique*  
 300  *$u \in W^{1,p}(\varpi, \Omega)/\mathbb{R}$  such that*

$$301 \quad \int_{\Omega} \nabla u \nabla \varphi = \int_{\Omega} \mathbf{f} \nabla \varphi, \quad \forall \varphi \in W^{1,p'}(\varpi, \Omega)$$

302 with the estimate

$$303 \quad \|\nabla u\|_{\mathbf{L}^p(\varpi,\Omega)} \lesssim \|\mathbf{f}\|_{\mathbf{L}^p(\varpi,\Omega)},$$

304 where the hidden constant depends on  $\Omega$ ,  $[\varpi]_{A_p}$  and  $p$ , but it is independent of  $\mathbf{f}$ .

305 *Proof.* All that is needed are the analogues of Theorems 2 and 4 to be able to  
 306 proceed as before. For that, we use [10, Theorem 3] and [15, Theorem 2], respectively.  $\square$

307 This immediately allows us to obtain a different Helmholtz decomposition, where  
 308 we exchange the boundary conditions from the space of gradients into the space of  
 309 solenoidal fields. Indeed, if given  $\varpi \in A_p(\Omega)$ , we define

$$310 \quad \mathbf{L}_{\sigma,D}^p(\varpi, \Omega) = \{\mathbf{v} \in \mathbf{L}^p(\varpi, \Omega) : \operatorname{div} \mathbf{v} = 0, \mathbf{v} \cdot \mathbf{n} = 0\},$$

311 where we denote by  $\mathbf{n}$  the outer normal to  $\Omega$  and

$$312 \quad \mathbf{G}_N^p(\varpi, \Omega) = \{\nabla v : v \in W^{1,p}(\varpi, \Omega)\},$$

313 then we can assert the following.

314 **COROLLARY 14** (weighted Helmholtz decomposition II). *In the setting of Theo-*  
 315 *rem 13 we have the following decomposition*

$$316 \quad (14) \quad \mathbf{L}^p(\varpi, \Omega) = \mathbf{L}_{\sigma,D}^p(\varpi, \Omega) \oplus \mathbf{G}_N^p(\varpi, \Omega).$$

317 *Proof.* Repeat the proof of Corollary 11 but using now Theorem 13.  $\square$

318 **4. The Stokes problem.** With techniques similar to the ones used to prove  
 319 Theorem 5 we can prove the well posedness of the Stokes problem (2) with singular  
 320 data  $\mathbf{F}$  and  $g$ . We begin by remarking that, owing to the boundary conditions on  $\mathbf{u}$ ,  
 321 we must necessarily have

$$322 \quad \int_{\Omega} g = 0.$$



365 Indeed, by using the cutoff function  $\psi_i$  and defining  $\mathbf{u}_i := \mathbf{u}\psi_i$  and  $\pi_i := \pi\psi_i$ , we  
 366 observe that  $(\mathbf{u}_i, \pi_i) \in \mathbf{W}_0^{1,p}(\varpi, \Omega_i) \times L^p(\varpi, \Omega_i)$  solve (15) with

$$367 \quad \int_{\Omega_i} \mathbf{F}_i \nabla \varphi = \int_{\Omega} \mathbf{F} \nabla(\varphi \psi_i) + \int_{\mathcal{G}} \mathbf{u} \otimes \nabla \psi_i \nabla \varphi + \int_{\mathcal{G}} \mathbf{u} \operatorname{div}(\nabla \psi_i \otimes \varphi) + \int_{\mathcal{G}} \pi \varphi \nabla \psi_i,$$

$$368 \quad \int_{\Omega_i} g_i q = \int_{\Omega} g \psi_i q + \int_{\mathcal{G}} \mathbf{u} \nabla \psi_i q,$$
 369

370 where  $\varphi \in \mathbf{C}_0^\infty(\Omega_i)$  and  $q \in C_0^\infty(\Omega_i)$ . Consequently, the estimates of Theorem 15  
 371 yield that

$$372 \quad \|\nabla \mathbf{u}_i\|_{\mathbf{L}^p(\varpi, \Omega_i)} + \|\pi_i\|_{L^p(\varpi, \Omega_i)} \lesssim \|\mathbf{F}_i\|_{\mathbf{L}^p(\varpi, \Omega_i)} + \|g_i\|_{L^p(\varpi, \Omega_i)}$$

373 with

$$374 \quad \|g_i\|_{L^p(\varpi, \Omega_i)} = \sup_{0 \neq q \in C_0^\infty(\Omega_i)} \frac{\int_{\Omega_i} g_i q}{\|q\|_{L^{p'}(\varpi', \Omega_i)}} \lesssim \|g\|_{L^p(\varpi, \Omega)} + \|\mathbf{u}\|_{\mathbf{L}^p(\mathcal{G})}$$

375 and

$$376 \quad \|\mathbf{F}_i\|_{\mathbf{L}^p(\varpi, \Omega_i)} \lesssim \|\mathbf{F}\|_{\mathbf{L}^p(\varpi, \Omega)} + \|\mathbf{u}\|_{\mathbf{L}^p(\mathcal{G})} + \sup_{0 \neq \varphi \in \mathbf{C}_0^\infty(\Omega_i)} \frac{\int_{\mathcal{G}} \pi \varphi \nabla \psi_i}{\|\nabla \varphi\|_{\mathbf{L}^{p'}(\varpi', \Omega_i)}}$$

$$377 \quad \lesssim \|\mathbf{F}\|_{\mathbf{L}^p(\varpi, \Omega)} + \|\mathbf{u}\|_{\mathbf{L}^p(\mathcal{G})} + \|\pi\|_{W^{-1,p}(\varpi, \Omega_i)}.$$

379 We now use the cutoff function  $\psi_\partial$  to define the functions  $\mathbf{u}_\partial = \mathbf{u}\psi_\partial \in \mathbf{W}^{1,p}(\mathcal{G})$   
 380 and  $\pi_\partial = \pi\psi_\partial \in L^p(\mathcal{G})$ . A similar calculation, together with Theorem 16 gives then  
 381 the desired bound for  $(\mathbf{u}_\partial, \pi_\partial)$  and, thus, (17).

- 382 • *Uniqueness:* We now prove that  $\mathbf{F} = \mathbf{0}$  and  $g = 0$  imply  $\mathbf{u} = \mathbf{0}$  and  $\pi = 0$ .  
 383 The argument is similar to Lemma 7. We first observe that, by [11, Theorem  
 384 IV.4.2] we have  $(\mathbf{u}_i, \pi_i) \in \mathbf{W}^{2,r}(\Omega_i) \times W^{1,r}(\Omega_i) \hookrightarrow \mathbf{W}^{1,2}(\Omega_i) \times L^2(\Omega_i)$ . In addition  
 385  $(\mathbf{u}_\partial, \pi_\partial) \in \mathbf{W}^{1,p}(\varpi, \mathcal{G}) \times L^p(\varpi, \mathcal{G}) \hookrightarrow \mathbf{W}^{1,2}(\mathcal{G}) \times L^2(\mathcal{G})$ .
- 386 • *A priori estimate (16):* This is, once again, proved by contradiction. We assume  
 387 (16) is false so that exist sequences

$$388 \quad (\mathbf{u}_k, \pi_k) \in \mathbf{W}_0^{1,p}(\varpi, \Omega) \times L^p(\varpi, \Omega)/\mathbb{R}, \quad (\mathbf{F}_k, g_k) \in \mathbf{L}^p(\varpi, \Omega) \times L^p(\varpi, \Omega)/\mathbb{R}$$

389 such that  $\|\nabla \mathbf{u}_k\|_{\mathbf{L}^p(\varpi, \Omega)} + \|\pi_k\|_{L^p(\varpi, \Omega)} = 1$  but that  $\|\mathbf{F}_k\|_{\mathbf{L}^p(\varpi, \Omega)} + \|g_k\|_{L^p(\varpi, \Omega)} \rightarrow 0$ .  
 390 Extracting weakly convergent subsequences and using uniqueness we conclude that  
 391 the limits must be  $\mathbf{u} = \mathbf{0}$  and  $\pi = 0$ . However, by compactness and (17)

$$392 \quad 1 = \|\nabla \mathbf{u}_k\|_{\mathbf{L}^p(\varpi, \Omega)} + \|\pi_k\|_{L^p(\varpi, \Omega)}$$

$$393 \quad \lesssim \|\mathbf{F}_k\|_{\mathbf{L}^p(\varpi, \Omega)} + \|g_k\|_{L^p(\varpi, \Omega)} + \|\mathbf{u}_k\|_{\mathbf{L}^p(\mathcal{G})} + \|\pi_k\|_{W^{-1,p}(\varpi, \Omega_i)} + \|\pi_k\|_{W^{-1,p}(\mathcal{G})}$$

$$394 \quad \rightarrow 0, \quad k \uparrow \infty,$$
 395

396 which is a contradiction.

- 397 • *Existence:* Finally, we construct a solution by approximation. For that, it suffices  
 398 to invoke the interior regularity of [11, Theorem IV.4.2].

399 This concludes the proof.  $\square$

400 **Acknowledgements.** AJS would like to thank his colleague Tadele Mengesha  
 401 for fruitful discussions. The authors would like to thank the anonymous reviewer for  
 402 carefully reading the manuscript and providing insightful suggestions.

403

## REFERENCES

- 404 [1] J. P. AGNELLI, E. M. GARAU, AND P. MORIN, *A posteriori error estimates for elliptic problems*  
 405 *with Dirac measure terms in weighted spaces*, ESAIM Math. Model. Numer. Anal., 48  
 406 (2014), pp. 1557–1581, <http://dx.doi.org/10.1051/m2an/2014010>.
- 407 [2] H. AIMAR, M. CARENA, R. DURÁN, AND M. TOSCHI, *Powers of distances to lower dimensional*  
 408 *sets as Muckenhoupt weights*, Acta Math. Hungar., 143 (2014), pp. 119–137, <http://dx.doi.org/10.1007/s10474-014-0389-1>.
- 409 [3] A. BONITO, R. A. DEVORE, AND R. H. NOCHETTO, *Adaptive finite element methods for elliptic*  
 410 *problems with discontinuous coefficients*, SIAM J. Numer. Anal., 51 (2013), pp. 3106–3134,  
 411 <http://dx.doi.org/10.1137/130905757>.
- 412 [4] M. BULÍČEK, L. DIENING, AND S. SCHWARZACHER, *Existence, uniqueness and optimal regularity*  
 413 *results for very weak solutions to nonlinear elliptic systems*, Anal. PDE, 9 (2016), pp. 1115–  
 414 1151, <https://doi.org/10.2140/apde.2016.9.1115>.
- 415 [5] M. BULÍČEK, J. BURČÁK, AND S. SCHWARZACHER, *A unified theory for some non-Newtonian*  
 416 *fluids under singular forcing*, SIAM J. Math. Anal., 48 (2016), pp. 4241–4267, <https://doi.org/10.1137/16M1073881>.
- 417 [6] C. D’ANGELO, *Finite element approximation of elliptic problems with Dirac measure terms in*  
 418 *weighted spaces: applications to one- and three-dimensional coupled problems*, SIAM J.  
 419 Numer. Anal., 50 (2012), pp. 194–215, <https://doi.org/10.1137/100813853>.
- 420 [7] J. DUOANDIKOETXEA, *Fourier analysis*, vol. 29 of Graduate Studies in Mathematics, American  
 421 Mathematical Society, Providence, RI, 2001. Translated and revised from the 1995 Spanish  
 422 original by David Cruz-Uribe.
- 423 [8] R. DURÁN AND I. OJEA, *A weighted setting for the Poisson problem with singular sources*. In  
 424 preparation.
- 425 [9] R. FARWIG AND H. SOHR, *Weighted  $L^q$ -theory for the Stokes resolvent in exterior domains*, J.  
 426 Math. Soc. Japan, 49 (1997), pp. 251–288, <https://doi.org/10.2969/jmsj/04920251>.
- 427 [10] A. FRÖHLICH, *The Helmholtz decomposition of weighted  $L^q$ -spaces for Muckenhoupt weights*,  
 428 Ann. Univ. Ferrara Sez. VII (N.S.), 46 (2000), pp. 11–19. Navier-Stokes equations and  
 429 related nonlinear problems (Ferrara, 1999).
- 430 [11] G. P. GALDI, *An introduction to the mathematical theory of the Navier-Stokes equations*,  
 431 Springer Monographs in Mathematics, Springer, New York, second ed., 2011, <https://doi.org/10.1007/978-0-387-09620-9>. Steady-state problems.
- 432 [12] D. GILBARG AND N. S. TRUDINGER, *Elliptic partial differential equations of second order*,  
 433 vol. 224 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles  
 434 of Mathematical Sciences], Springer-Verlag, Berlin, second ed., 1983, <https://doi.org/10.1007/978-3-642-61798-0>.
- 435 [13] V. GOL’DSHTEIN AND A. UKHLOV, *Weighted Sobolev spaces and embedding theorems*,  
 436 Trans. Amer. Math. Soc., 361 (2009), pp. 3829–3850, <http://dx.doi.org/10.1090/S0002-9947-09-04615-7>.
- 437 [14] Q. HAN AND F. LIN, *Elliptic partial differential equations*, vol. 1 of Courant Lecture Notes in  
 438 Mathematics, Courant Institute of Mathematical Sciences, New York; American Mathe-  
 439 matical Society, Providence, RI, second ed., 2011.
- 440 [15] D. JERISON AND C. E. KENIG, *The functional calculus for the Laplacian on Lipschitz domains*,  
 441 in Journées “Équations aux Dérivées Partielles” (Saint Jean de Monts, 1989), École Poly-  
 442 tech., Palaiseau, 1989, pp. Exp. No. IV, 10.
- 443 [16] D. JERISON AND C. E. KENIG, *The inhomogeneous Dirichlet problem in Lipschitz domains*, J.  
 444 Funct. Anal., 130 (1995), pp. 161–219, <https://doi.org/10.1006/jfan.1995.1067>.
- 445 [17] V. A. KOZLOV, V. G. MAZ’YA, AND J. ROSSMANN, *Elliptic boundary value problems in domains*  
 446 *with point singularities*, American Mathematical Society, Providence, Rhode Island, USA,  
 447 1997.
- 448 [18] N. G. MEYERS, *An  $L^p$ -estimate for the gradient of solutions of second order elliptic divergence*  
 449 *equations*, Ann. Scuola Norm. Sup. Pisa (3), 17 (1963), pp. 189–206.
- 450 [19] M. MITREA AND M. WRIGHT, *Boundary value problems for the Stokes system in arbitrary*  
 451 *Lipschitz domains*, Astérisque, (2012), pp. viii+241.
- 452 [20] B. MUCKENHOUT, *Weighted norm inequalities for the Hardy maximal function*, Trans. Amer.  
 453 Math. Soc., 165 (1972), pp. 207–226, <https://doi.org/10.2307/1995882>.

- 459 [21] R. H. NOCHETTO, E. OTÁROLA, AND A. J. SALGADO, *Piecewise polynomial interpolation in*  
460 *Muckenhoupt weighted Sobolev spaces and applications*, Numer. Math., 132 (2016), pp. 85–  
461 130, <http://dx.doi.org/10.1007/s00211-015-0709-6>.
- 462 [22] B. O. TURESSON, *Nonlinear potential theory and weighted Sobolev spaces*, vol. 1736 of Lec-  
463 ture Notes in Mathematics, Springer-Verlag, Berlin, 2000, [http://dx.doi.org/10.1007/](http://dx.doi.org/10.1007/BFb0103908)  
464 [BFb0103908](http://dx.doi.org/10.1007/BFb0103908).